

WASTE STREAM	2C313	Stainless Steel (Reactor) ILW
---------------------	--------------	--------------------------------------

SITE Chapelcross
SITE OWNER Nuclear Decommissioning Authority
WASTE CUSTODIAN Magnox Limited
WASTE TYPE ILW

WASTE VOLUMES

		Reported
Stocks:	At 1.4.2019.....	0 m ³
Future arisings -	1.4.2089 - 31.3.2095.....	80.0 m ³
Total future arisings:		80.0 m ³
Total waste volume:		80.0 m ³

Comment on volumes: For inventory purposes the arisings are assumed to arise at a uniform rate over six years. Final Dismantling & Site Clearance is assumed to commence in 2085 with reactor dismantling commencing in 2089 and lasting for 6 years. The volumes and radioactivity have been calculated for 85 years after reactor shutdown, i.e. 2089.

Uncertainty factors on volumes: Stock (upper): x Arisings (upper) x 1.2
 Stock (lower): x Arisings (lower) x 0.8

WASTE SOURCE Stainless steel items from reactor dismantling.

PHYSICAL CHARACTERISTICS

General description: A variety of stainless steel items.
 Physical components (%vol): Stainless steel items (100%).
 Sealed sources: -
 Bulk density (t/m³): 1.4
 Comment on density: The density is of the waste as cut for packaging.

CHEMICAL COMPOSITION

General description and components (%wt): Stainless steel items (100%).
 Chemical state: Neutral
 Chemical form of radionuclides: H-3: The tritium content is insignificant.
 C-14: The carbon 14 is incorporated in the steel. There may also be some contamination as graphite.
 Se-79: The selenium content is insignificant.
 Tc-99: The chemical form of technetium has not been determined.
 Ra: The radium content is insignificant.
 Th: The thorium content is insignificant.
 U: The uranium content is insignificant.
 Np: The neptunium content is insignificant.
 Pu: The plutonium content is insignificant.

Metals and alloys (%wt): Items have been cut for packaging. Thicknesses are likely to vary from a few mm to about 25 mm.

Stainless steel.....	100.0	Stainless steel types are BS970-EN56 and BS970-EN58B.
Other ferrous metals.....	0	
Iron.....		
Aluminium.....	0	
Beryllium.....		
Cobalt.....		
Copper.....	0	
Lead.....	0	
Magnox/Magnesium.....	0	

WASTE STREAM	2C313	Stainless Steel (Reactor) ILW
---------------------	--------------	--------------------------------------

	Nickel.....	
	Titanium.....	
	Uranium.....	
	Zinc.....	0
	Zircaloy/Zirconium.....	0
	Other metals.....	0
		There are no "other" metals.
Organics (%wt):	None expected. No halogenated plastics or rubbers will be present.	
	Total cellulosics.....	0
	Paper, cotton.....	0
	Wood.....	0
	Halogenated plastics	0
	Total non-halogenated plastics.....	0
	Condensation polymers.....	0
	Others.....	0
	Organic ion exchange materials....	0
	Total rubber.....	0
	Halogenated rubber	0
	Non-halogenated rubber.....	0
	Hydrocarbons.....	
	Oil or grease	
	Fuel.....	
	Asphalt/Tarmac (cont.coal tar)...	
	Asphalt/Tarmac (no coal tar)....	
	Bitumen.....	
	Others.....	
	Other organics.....	0
Other materials (%wt):	Some graphite dust may be associated with reactor materials.	
	Inorganic ion exchange materials.	0
	Inorganic sludges and flocs.....	0
	Soil.....	0
	Brick/Stone/Rubble.....	0
	Cementitious material.....	0
	Sand.....	
	Glass/Ceramics.....	0
	Graphite.....	TR
	Desiccants/Catalysts.....	
	Asbestos.....	0
	Non/low friable.....	
	Moderately friable.....	
	Highly friable.....	
	Free aqueous liquids.....	0
	Free non-aqueous liquids.....	0
	Powder/Ash.....	0

WASTE STREAM	2C313	Stainless Steel (Reactor) ILW
---------------------	--------------	--------------------------------------

Inorganic anions (%wt): Trace quantities of chloride may be present.

Fluoride.....	0
Chloride.....	TR
Iodide.....	0
Cyanide.....	0
Carbonate.....	0
Nitrate.....	0
Nitrite.....	0
Phosphate.....	0
Sulphate.....	0
Sulphide.....	0

Materials of interest for waste acceptance criteria: No materials likely to pose a fire or other non-radiological hazard have been identified.

Combustible metals.....	0
Low flash point liquids.....	0
Explosive materials.....	0
Phosphorus.....	0
Hydrides.....	0
Biological etc. materials.....	0
Biodegradable materials.....	
Putrescible wastes.....	0
Non-putrescible wastes.....	
Corrosive materials.....	0
Pyrophoric materials.....	0
Generating toxic gases.....	0
Reacting with water.....	0
Active particles.....	
Soluble solids as bulk chemical compounds.....	

Hazardous substances / non hazardous pollutants: None expected.

Acrylamide.....	
Benzene.....	
Chlorinated solvents.....	
Formaldehyde.....	
Organometallics.....	
Phenol.....	
Styrene.....	
Tri-butyl phosphate.....	
Other organophosphates.....	
Vinyl chloride.....	
Arsenic.....	
Barium.....	

WASTE STREAM 2C313 Stainless Steel (Reactor) ILW

Boron.....
 Cadmium.....
 Caesium.....
 Selenium.....
 Chromium.....
 Molybdenum.....
 Thallium.....
 Tin.....
 Vanadium.....
 Mercury compounds.....
 Others.....
 Electronic Electrical Equipment (EEE)
 EEE Type 1.....
 EEE Type 2.....
 EEE Type 3.....
 EEE Type 4.....
 EEE Type 5.....

Complexing agents (%wt):

EDTA.....
 DPTA.....
 NTA.....
 Polycarboxylic acids.....
 Other organic complexants.....
 Total complexing agents..... TR

PACKAGING AND CONDITIONING

Conditioning method: The waste is not expected to be supercompacted. It will be placed in baskets in the waste packages, and is now assumed to be encapsulated.

Plant Name: -

Location: Chapelcross Power Station

Plant startup date: 2089

Total capacity (m³/y incoming waste): ~5000.0

Target start date for packaging this stream: 2089

Throughput for this stream (m³/y incoming waste): ~11.5

Other information: -

Likely container type:	Container	Waste packaged (%vol)	Waste loading (m ³)	Payload (m ³)	Number of packages
	4m box (200mm concrete shielding)	100.0	9.4	10.9	9

WASTE STREAM 2C313 Stainless Steel (Reactor) ILW

Likely container type comment: The waste is assumed to be in baskets in the waste package so the occupied volume in the package is greater than the original waste volume.

Range in container waste volume: Not yet determined. No significant variability is expected.

Other information on containers: The container material is expected to be stainless steel. Container choice may be influenced by Transport Regulations at the time of final site clearance.

Likely conditioning matrix: Blast Furnace Slag / Ordinary Portland Cement

Other information: The waste is now assumed to be encapsulated.

Conditioned density (t/m³): 3.0

Conditioned density comment: The conditioned waste density assumes the waste will be encapsulated.

Other information on conditioning: The waste will be in baskets placed in the waste packages. Baskets of different Final Dismantling & Site Clearance ILW wastes may be in the same waste package. Should encapsulation not be required the density of the conditioned waste product would be about 1.2 t/m³.

Opportunities for alternative disposal routing: No

Treatment	Stream volume (%)	Comment
-	-	-

RADIOACTIVITY

Source: Activation of the stainless steel and impurities.

Uncertainty: The values quoted were derived by calculation from available material specification and are indicative of the activities that are expected. The major source of uncertainty is the impurity levels.

Definition of total alpha and total beta/gamma: Total beta/gamma is defined as the sum of the listed activities of all nuclides other than alpha emitters. All alpha emitter activities are insignificant.

Measurement of radioactivities: Specific Activities have been estimated using a neutron activation calculation.

Other information: The activities quoted are those at 85 years after reactor shutdown, i.e. in 2089. There may be some contamination by Cs137.

WASTE STREAM 2C313 Stainless Steel (Reactor) ILW

Nuclide	Mean radioactivity, TBq/m ³				Nuclide	Mean radioactivity, TBq/m ³			
	Waste at 1.4.2019	Bands and Code	Future arisings	Bands and Code		Waste at 1.4.2019	Bands and Code	Future arisings	Bands and Code
H 3				8	Gd 153				8
Be 10				8	Ho 163				8
C 14			8.62E-02	C C 2	Ho 166m				8
Na 22				8	Tm 170				8
Al 26				8	Tm 171				8
Cl 36			6.67E-06	C C 2	Lu 174				8
Ar 39				8	Lu 176				8
Ar 42				8	Hf 178n				8
K 40				8	Hf 182				8
Ca 41				8	Pt 193				8
Mn 53				8	Tl 204		3.23E-07	C C 2	8
Mn 54				8	Pb 205				8
Fe 55			1.4E-07	C C 2	Pb 210				8
Co 60			2.25E-03	C C 2	Bi 208				8
Ni 59			2.17E-01	C C 2	Bi 210m				8
Ni 63			1.3E+01	C C 2	Po 210				8
Zn 65				8	Ra 223				8
Se 79				8	Ra 225				8
Kr 81				8	Ra 226				8
Kr 85				8	Ra 228				8
Rb 87				8	Ac 227				8
Sr 90				8	Th 227				8
Zr 93				8	Th 228				8
Nb 91				8	Th 229				8
Nb 92			4.06E-09	C C 2	Th 230				8
Nb 93m				8	Th 232				8
Nb 94			5.41E-04	C C 2	Th 234				8
Mo 93			6.69E-03	C C 2	Pa 231				8
Tc 97				8	Pa 233				8
Tc 99			6.49E-05	C C 2	U 232				8
Ru 106				8	U 233				8
Pd 107				8	U 234				8
Ag 108m			2.98E-03	C C 2	U 235				8
Ag 110m				8	U 236				8
Cd 109				8	U 238				8
Cd 113m				8	Np 237				8
Sn 119m				8	Pu 236				8
Sn 121m				8	Pu 238				8
Sn 123				8	Pu 239				8
Sn 126				8	Pu 240				8
Sb 125				8	Pu 241				8
Sb 126				8	Pu 242				8
Te 125m				8	Am 241				8
Te 127m				8	Am 242m				8
I 129				8	Am 243				8
Cs 134				8	Cm 242				8
Cs 135				8	Cm 243				8
Cs 137				6	Cm 244				8
Ba 133				8	Cm 245				8
La 137				8	Cm 246				8
La 138				8	Cm 248				8
Ce 144				8	Cf 249				8
Pm 145				8	Cf 250				8
Pm 147				8	Cf 251				8
Sm 147				8	Cf 252				8
Sm 151				8	Other a				
Eu 152				8	Other b/g				
Eu 154				8	Total a	0	0		
Eu 155				8	Total b/g	0	1.33E+01	C C 2	

Bands (Upper and Lower)

- A a factor of 1.5
- B a factor of 3
- C a factor of 10
- D a factor of 100
- E a factor of 1000

Note: Bands quantify uncertainty in mean radioactivity.

Code

- 1 Measured activity
- 2 Derived activity (best estimate)
- 3 Derived activity (upper limit)
- 4 Not present
- 5 Present but not significant
- 6 Likely to be present but not assessed
- 7 Present in significant quantities but not determined
- 8 Not expected to be present in significant quantity