SITE Chapelcross

SITE OWNER Nuclear Decommissioning Authority

WASTE CUSTODIAN Magnox Limited

WASTE TYPE ILW

Is the waste subject to

Scottish Policy:

Yes

WASTE VOLUMES

Reported

Total future arisings: 3647.0 m³
Total waste volume: 3647.0 m³

Comment on volumes: For inventory purposes the arisings are assumed to arise at a uniform rate over 6 years.

Final Dismantling & Site Clearance is assumed to commence in 2085 with reactor dismantling commencing in 2089 and lasting for 6 years. The volumes and radioactivity

have been calculated for 85 years after reactor shutdown, i.e. 2089

Uncertainty factors on Stock (upper): x Arisings (upper) x 1.2 volumes: Stock (lower): x Arisings (lower) x 0.8

WASTE SOURCE Moderator and reflector graphite from reactor dismantling.

PHYSICAL CHARACTERISTICS

General description: Graphite blocks and other graphite components.

Physical components (%vol): Graphite (100%).

Sealed sources: The waste does not contain sealed sources.

Bulk density (t/m³): 1.25

Comment on density: Density estimate based upon the assumed packing efficiency of the waste. The density is

the effective density for packaging assuming 90% of the graphite is in blocks and 10% is

rubble.

CHEMICAL COMPOSITION

General description and

components (%wt):

Graphite and possibly traces of ferrous metals.

Chemical state: Neutral

Chemical form of H-3: Tritium may be chemically bound with the graphite.

radionuclides: C-14: The carbon 14 will be present as graphite.

Se-79: The selenium content is insignificant. Tc-99: The technetium content is insignificant. Ra: The radium content is insignificant. Th: The thorium content is insignificant.

U: There may be traces of uranium as metal or oxide.

Np: The neptunium content is insignificant.

Pu: There may be traces of plutonium as metal or oxide.

Metals and alloys (%wt): There are no metallic items present.

(%wt) Type(s) / Grade(s) with proportions % of total C14

activity

Stainless steel...... 0

Other ferrous metals...... TR There may be trace contamination

by ferrous metals.

Iron.....

Aluminium...... 0

Beryllium.....
Cobalt.....

	Copper	0		
	Lead	0		
	Magnox/Magnesium	0		
	Nickel			
	Titanium			
	Uranium			
	Zinc	0		
	Zircaloy/Zirconium	. 0		
	Other metals	0	There are no "other" metals present.	
Organio	cs (%wt): None expected. No	halogenat	ted plastics or rubbers will be present.	
		(%wt)	Type(s) and comment	% of total C14
	Total cellulosics	0		activity
	Paper, cotton	0		
	Wood	0		
	Halogenated plastics	0		
	Total non-halogenated plastics	0		
	Condensation polymers	0		
	Others	0		
	Organic ion exchange materials	0		
	Total rubber	0		
	Halogenated rubber	0		
	Non-halogenated rubber	0		
	Hydrocarbons	Ü		
	Oil or grease			
	Fuel			
	Asphalt/Tarmac (cont.coal tar) Asphalt/Tarmac (no coal tar)			
	Bitumen			
	Others			
	Other organics	0		
	•	0		
Other n	naterials (%wt): -			
		(%wt)	Type(s) and comment	% of total C14
	la cracació ion evolución a motoriole	0		activity
	Inorganic ion exchange materials	0		
	Inorganic sludges and flocs	0		
	Soil	0		
	Brick/Stone/Rubble	0		
	Cementitious material	0		
	Sand Glass/Ceramics	0		
		0		100.0
	Graphite	100.0		100.0
	Desiccants/Catalysts			

Asbestos	0	
Non/low friable		
Moderately friable		
Highly friable		
Free aqueous liquids	0	
Free non-aqueous liquids	0	
Powder/Ash	0	
Inorganic anions (%wt): None of the inorgative concentration		isted in the table is expected to be present at greater than
	(%wt)	Type(s) and comment
Fluoride	TR	Detected at trace levels in inactive graphite material.
Chloride	TR	
lodide	0	
Cyanide	0	
Carbonate	TR	
Nitrate	TR	
Nitrite	TR	
Phosphate	TR	Detected at trace levels in inactive graphite material.
Sulphate	TR	Detected at trace levels in inactive graphite material.
Sulphide	0	
		re or other non-radiological hazard have been identified. sk; it is difficult but not impossible to ignite.
	(%wt)	Type(s) and comment
Combustible metals	0	
Low flash point liquids	0	
Explosive materials	0	
Phosphorus	TR	Detected at trace levels in inactive graphite material.
Hydrides	0	
Biological etc. materials	0	
Biodegradable materials		
Putrescible wastes	0	
Non-putrescible wastes		
Corrosive materials	0	
Pyrophoric materials	0	
Generating toxic gases	0	
Reacting with water	0	
Higher activity particles		
Soluble solids as bulk chemical compounds		

Hazardous substances /	
non hazardous pollutants:	

Complexing

None expected

	(%wt)	Type(s) and comment
Acrylamide		
Benzene		
Chlorinated solvents		
Formaldehyde		
Organometallics		
Phenol		
Styrene		
Tri-butyl phosphate		
Other organophosphates		
Vinyl chloride		
Arsenic	TR	Detected at trace levels in inactive graphite material.
Barium		
Boron		
Boron (in Boral)		
Boron (non-Boral)		
Cadmium		
Caesium		
Selenium		
Chromium		
Molybdenum		
Thallium	TR	Detected at trace levels in inactive graphite material.
Tin		
Vanadium		
Mercury compounds		
Others	TR	Gallium, germanium and rubidium detected at trace levels in inactive graphite material.
Electronic Electrical Equipment (EEE)		
EEE Type 1		
EEE Type 2		
EEE Type 3		
EEE Type 4		
EEE Type 5		
agents (%wt): Yes		
	(%wt)	Type(s) and comment
EDTA		
DPTA		
NTA		
Polycarboxylic acids		
Other organic complexants		

Total complexing agents..... TR

Potential for the waste to contain discrete items:

Yes. Graphite Bricks/Tiles assumed to be Dls. Bricks assumed drummed (ungrouted) so assumed Bricks are DIs; If grouted, Drum is also a DI. "Rubble" pieces assumed drummed (ungrouted) assumed NOT DIs; If grouted, Drum is a

PACKAGING AND CONDITIONING

Conditioning method: The waste is not expected to be supercompacted. It will be placed in baskets in the

waste packages, and is now assumed to be encapsulated.

Plant Name:

Location: Chapelcross Power Station

2089 Plant startup date:

Total capacity

(m³/y incoming waste):

2089

Target start date for packaging this stream:

Throughput for this stream (m³/y incoming waste):

~521.0

Other information:

Likely container type:

Container	Waste packaged (%vol)	Waste loading (m³)	Payload (m³)	Number of packages
4m box (no shielding)	100.0	~16.2	18.9	226

Likely container type

comment:

The waste is assumed to be in baskets in the waste package so the occupied volume in

the package is greater than the original waste volume.

Range in container waste

volume:

Not yet determined. No significant variability is expected.

Other information on

containers:

The container material is expected to be stainless steel. Container choice may be

influenced by Transport Regulations at the time of final site clearance.

Likely conditioning matrix:

Other information:

Blast Furnace Slag / Ordinary Portland Cement

It is now assumed that the waste will be encapsulated. The matrix could be BFS/OPC.

Conditioned density (t/m³): Conditioned density

comment:

The conditioned waste density assumes that the waste will be encapsulated.

Other information on

conditioning:

The waste will be in baskets placed in the waste packages. Baskets of different Final Dismantling & Site Clearance ILW wastes may be in the same waste package. Should encapsulation not be required the density of the conditioned waste product would be about 1.1 t/m3. Data have been presented as if the waste will be placed in a container with other

ILW.

Opportunities for alternative disposal routing:

Baseline Management Route	Opportunity Management Route	Stream volume (%)	Estimated Date that Opportunity	Opportunity Confidence	Comment
Management Route	Management Route	volunio (70)	will be realised	Communico	

RADIOACTIVITY

Source: Activation of the graphite and impurities.

Uncertainty: The values quoted were derived by calculation from available material specification and

are indicative of the activities that are expected. The major source of uncertainty is the

impurity levels.

Definition of total alpha and total beta/gamma:

Where totals are shown on the table of radionuclide activities they are the sums of the listed alpha or beta/gamma emitting radionuclides plus 'other alpha' or 'other beta/gamma'.

Measurement of radioactivities:

The specific activities have been estimated using a neutron activation calculation. With additional data from newly calculated inventories including 100 ppb U precursor as per M/EF/GEN/EAN/0008/20. Additionally 440ppb of fuel contamination was included in CHA streams because of the known fuel fire event in 1967 - although this had a relatively small

impact.

Other information: There may be some contamination by Cs137. The activities quoted are those at 85 years

after reactor shutdown, i.e. in 2089. Fission of trace uranium impurity in the graphite may

result in some fission product and actinide activity.

WASTE STREAM Graphite ILW 2C311

	Mean radioactivity, TBq/m³			Mean radioactivity, TBq/m³					
Nuclide	Waste at 1.4.2022	Bands and Code	Future arisings	Bands and Code	Nuclide	Waste at 1.4.2022	Bands and Code	Future arisings	Bands and Code
H 3			2.76E-03	CC 2	Gd 153				8
Be 10				8	Ho 163				8
C 14			1.28E-01	BB 2	Ho 166m			1.99E-05	CC 2
Na 22				8	Tm 170				8
AI 26				8	Tm 171				8
CI 36			2.94E-04	CC 2	Lu 174				8
Ar 39				8	Lu 176				8
Ar 42				8	Hf 178n				8
K 40				8	Hf 182				8
Ca 41			3.44E-04	CC 2	Pt 193				8
Mn 53				8	TI 204				8
Mn 54				8	Pb 205				8
Fe 55				8	Pb 210				8
Co 60			3.3E-06	CC 2	Bi 208				8
Ni 59			6.25E-05	CC 2	Bi 210m				8
Ni 63			4.76E-03	CC 2	Po 210				8
Zn 65	İ			8	Ra 223				8
Se 79			1.56E-09	CC 2	Ra 225				8
Kr 81				8	Ra 226				8
Kr 85			9.72E-07	CC 2	Ra 228				8
Rb 87				8	Ac 227				8
Sr 90			2.98E-04	CC 2	Th 227				8
Zr 93			8.36E-08	CC 2	Th 228				8
Nb 91				8	Th 229				8
Nb 92				8	Th 230				8
Nb 93m			7.6E-08	CC 2	Th 232				8
Nb 94			1.82E-05	CC 2	Th 234				8
Mo 93				8	Pa 231				8
Tc 97				8	Pa 233			1.05E-09	CC 2
Tc 99			5.3E-07	CC 2	U 232				8
Ru 106	İ		0.02	8	U 233				8
Pd 107			7.62E-09	CC 2	U 234			1.23E-08	CC 2
Ag 108m			8.72E-07	CC 2	U 235				8
Ag 110m	İ		0.722 07	8	U 236				8
Cd 109				8	U 238			7.4E-09	CC 2
Cd 113m				8	Np 237			1.28E-09	CC 2
Sn 119m				8	Pu 236				8
Sn 121m			1.83E-04	CC 2	Pu 238			1.64E-05	CC 2
Sn 123				8	Pu 239	ĺ		4.12E-06	CC 2
Sn 126			3.13E-08	CC 2	Pu 240			1.76E-05	CC 2
Sb 125			002 00	8	Pu 241			2.31E-05	CC 2
Sb 126			4.17E-09	CC 2	Pu 242			2.16E-07	CC 2
Te 125m			1.172 00	8	Am 241			3.69E-05	CC 2
Te 127m	[8	Am 242m	İ		4.96E-08	CC 2
I 129	[8	Am 243			5.43E-06	CC 2
Cs 134				8	Cm 242			3.45E-08	CC 2
Cs 134	[2.45E-08	CC 2	Cm 243			5.32E-08	CC 2
Cs 133	1		5.78E-04	CC 2	Cm 244			1.58E-04	CC 2
Ba 133	[5.76E-04 5.94E-07	CC 2	Cm 245			3.21E-07	CC 2
La 137	1		5.5 4 L-67	8	Cm 246			2.31E-06	CC 2
La 137	[8	Cm 248	i		2.012 00	8
	[8	Cf 249			7.57E-09	CC 2
Ce 144 Pm 145			6.88E-08	CC 2	Cf 249 Cf 250			1.23E-09	CC 2
Pm 145			0.00⊏-08			l I		1.236-08	
Pm 147	[8	Cf 251				8 8
Sm 147			2 44 5 00	8	Cf 252				0
Sm 151	[3.11E-06	CC 2	Other a				
Eu 152	1		4 4 4 5 00	8	Other b/g	_ ا		2 44 = 04	CC 3
Eu 154			1.14E-06	CC 2	Total a	0		2.41E-04	CC 2
Eu 155	I		4.14E-09	CC 2	Total b/g	0	-	1.37E-01	CC 2

Bands (Upper and Lower)

A a factor of 1.5 B a factor of 3

C a factor of 10

D a factor of 100

E a factor of 1000

Note: Bands quantify uncertainty in mean radioactivity.

- Measured activity
 Derived activity (best estimate)
 Derived activity (upper limit)
- 4 Not present
- 5 Present but not significant
- 6 Likely to be present but not assessed
- 7 Present in significant quantities but not determined
- 8 Not expected to be present in significant quantity