

WASTE STREAM	2D08.1	MSSS Compartment 1
---------------------	---------------	---------------------------

SITE Sellafield

SITE OWNER Nuclear Decommissioning Authority

WASTE CUSTODIAN Sellafield Limited

WASTE TYPE ILW

Is the waste subject to Scottish Policy: No

WASTE VOLUMES

Stocks:	At 1.4.2022.....	Reported ~606.0 m ³
---------	------------------	-----------------------------------

Total future arisings:	0 m ³
------------------------	------------------

Total waste volume:	606.0 m ³
---------------------	----------------------

Comment on volumes: There are no further arisings to this waste stream. Waste volume includes swarf/sludge, miscellaneous beta/gamma and aggregate. Waste volumes are estimates derived from known fuel decanning campaigns and storage volumes.

Uncertainty factors on volumes:	Stock (upper):	x 1.5	Arisings (upper)	x
	Stock (lower):	x 0.66	Arisings (lower)	x

WASTE SOURCE Swarf from Magnox decanning. Other miscellaneous beta/gamma wastes from Windscale and other sites.

PHYSICAL CHARACTERISTICS

General description: Magnox cladding, much of which has been converted to a magnesium hydroxide sludge. Some miscellaneous beta/gamma wastes (MBGW) are also present: these include tins/cans, filters, tools, pipework, pumps, valves, plastic items. The waste also includes some uranium most of which has corroded to oxides of uranium. A small volume of limestone aggregate is present at the bottom of the compartment. Swarf from Magnox decanning has reacted with the storage cover water to form corrosion products. Some degradation of miscellaneous components waste will have occurred.

Physical components (%wt): Magnox swarf (2.85%), Magnox sludge (35.08%), Al/Al(OH)₃ (1.10%), U/UO₂/U₃O₈ (1.71%), MBGW (scrap, filters, cans, etc.) (31.32%), aggregate (limestone) (1.16%), water (26.39%), others (0.38%). Note that interstitial liquor and cover liquor are included as part of the %wt (water).

Sealed sources: Not yet determined.

Bulk density (t/m³): 1.77

Comment on density: Bulk density is solids only - includes water content based on assumptions of draining characteristics. Bulk density including interstitial liquor = 1.77t/m³, bulk density with cover liquor and interstitial liquor = 1.75t/m³. Average decanning debris is ~1.6t/m³ and average MBGW is ~3.2t/m³.

CHEMICAL COMPOSITION

General description and components (%wt): Magnox swarf (2.85%), Magnox sludge (35.08%), Al/Al(OH)₃ (1.10%), U/UO₂/U₃O₈ (1.71%), MBGW (scrap, filters, cans, etc.) (31.32%), aggregate (limestone) (1.16%), water (26.39%), others (0.38%). Note that interstitial liquor and cover liquor are included as part of the %wt (water).

Chemical state: Neutral

Chemical form of radionuclides: H-3: Present in elemental and reacted forms.
 C-14: Present in elemental and reacted forms.
 Cl-36: Present in elemental and reacted forms.
 Se-79: Present in elemental and reacted forms.
 Tc-99: Present in elemental and reacted forms.
 I-129: Present in elemental and reacted forms.
 Ra: Present in elemental forms.
 Th: Present in elemental and reacted forms.
 U: Present in metallic and reacted forms (oxides and possibly hydride).
 Np: Present in elemental and reacted forms.
 Pu: Present in metallic and mixed oxide forms.

Metals and alloys (%wt): No significant quantities of sheet metal, although small quantities of fabricated items are present, e.g. paint tins, HEPA filters. Small pieces of fin Magnox metal present.

WASTE STREAM	2D08.1	MSSS Compartment 1
---------------------	---------------	---------------------------

	(%wt)	Type(s) / Grade(s) with proportions	% of total C14 activity
Stainless steel.....	~7.5	Mainly SS316 and SS304.	
Other ferrous metals.....	~16.5	Carbon steels.	
Iron.....	P		
Aluminium.....	~1.1		
Beryllium.....	NE		
Cobalt.....	P		
Copper.....	TR		
Lead.....	P		
Magnox/Magnesium.....	~2.8	Predominantly Magnox.	
Nickel.....	NE		
Titanium.....	NE		
Uranium.....	~0.63		
Zinc.....	TR		
Zircaloy/Zirconium.....	~0.03		
Other metals.....	~0.04		
Organics (%wt):	The organic content of the waste is low.		
	(%wt)	Type(s) and comment	% of total C14 activity
Total cellulose.....	~3.5		
Paper, cotton.....			
Wood.....			
Halogenated plastics	~0.65		
Total non-halogenated plastics.....	~0.60		
Condensation polymers.....			
Others.....			
Organic ion exchange materials....			
Total rubber.....	TR		
Halogenated rubber			
Non-halogenated rubber.....			
Hydrocarbons.....			
Oil or grease	<0.01		
Fuel.....	NE		
Asphalt/Tarmac (cont.coal tar)...	NE		
Asphalt/Tarmac (no coal tar)....	NE		
Bitumen.....	NE		
Others.....	NE		
Other organics.....	0.21		

Other materials (%wt): -

WASTE STREAM	2D08.1	MSSS Compartment 1
---------------------	---------------	---------------------------

	(%wt)	Type(s) and comment	% of total C14 activity
Inorganic ion exchange materials..	NE		
Inorganic sludges and flocs.....	~35.2		
Soil.....	NE		
Brick/Stone/Rubble.....	~1.1		
Cementitious material.....	NE		
Sand.....	NE		
Glass/Ceramics.....	~0.27		
Graphite.....	~0.66		
Desiccants/Catalysts.....	NE		
Asbestos.....	NE		
Non/low friable.....			
Moderately friable.....			
Highly friable.....			
Free aqueous liquids.....	~25.7	Includes interstitial liquor and cover liquor. Total waste volume 617 m3 with cover liquor.	
Free non-aqueous liquids.....	NE		
Powder/Ash.....	NE		

Inorganic anions (%wt): Sodium chloride was added during the filling of the compartments. Hydroxides and carbonates are present.

	(%wt)	Type(s) and comment
Fluoride.....	TR	
Chloride.....	~0.05	Alkali metal and alkaline earth chlorides.
Iodide.....	TR	
Cyanide.....	NE	
Carbonate.....	TR	
Nitrate.....	TR	
Nitrite.....	TR	
Phosphate.....	TR	
Sulphate.....	TR	
Sulphide.....	TR	

Materials of interest for waste acceptance criteria: Waste contains uranium, Magnox and traces of uranium hydride. Cellulosics are present as putrescible waste.

	(%wt)	Type(s) and comment
Combustible metals.....	0	
Low flash point liquids.....	0	
Explosive materials.....	0	
Phosphorus.....	0	
Hydrides.....	P	Trace, passivated.
Biological etc. materials.....	0	
Biodegradable materials.....	P	

WASTE STREAM	2D08.1	MSSS Compartment 1
---------------------	---------------	---------------------------

Putrescible wastes.....	P	Cellulosics.
Non-putrescible wastes.....	P	Organics - plastics and rubbers.
Corrosive materials.....	0	
Pyrophoric materials.....	0	
Generating toxic gases.....	0	
Reacting with water.....	P	<5%
Higher activity particles.....	P	Radioactive sludges and solids.
Soluble solids as bulk chemical compounds.....	P	Elemental and hydroxides, oxides and carbonates.

Hazardous substances / non hazardous pollutants: Lead (blocks and sheet) is present. Uranium and irradiated fuel components are present.

	(%wt)	Type(s) and comment
Acrylamide.....		
Benzene.....	0	
Chlorinated solvents.....		
Formaldehyde.....		
Organometallics.....		
Phenol.....	0	
Styrene.....		
Tri-butyl phosphate.....	0	
Other organophosphates.....		
Vinyl chloride.....	0	
Arsenic.....	0	
Barium.....		
Boron.....	NE	Traces - elemental and borates.
Boron (in Boral).....		
Boron (non-Boral).....		
Cadmium.....	0	
Caesium.....		
Selenium.....	0	
Chromium.....	P	In steels.
Molybdenum.....	P	In steels.
Thallium.....		
Tin.....	P	Traces.
Vanadium.....	P	In steels.
Mercury compounds.....		
Others.....	NE	
Electronic Electrical Equipment (EEE)		
EEE Type 1.....		
EEE Type 2.....		
EEE Type 3.....		
EEE Type 4.....		

WASTE STREAM 2D08.1 MSSS Compartment 1

EEE Type 5.....

Complexing agents (%wt): Not yet determined

(%wt) Type(s) and comment

EDTA.....

DPTA.....

NTA.....

Polycarboxylic acids.....

Other organic complexants.....

Total complexing agents.....

Potential for the waste to contain discrete items: Not yet determined.

PACKAGING AND CONDITIONING

Conditioning method: The waste will be retrieved from MSSS and stored in an unconditioned state for a period of 70 years. Following the period of storage it will be retrieved (from storage) and conditioned for ultimate disposal to the GDF.

Plant Name: TBD

Location: Sellafield

Plant startup date: 2070-2090

Total capacity (m³/y incoming waste): -

Target start date for packaging this stream: -

Throughput for this stream (m³/y incoming waste): -

Other information: The plant has not yet been designed. Hence, throughputs & capacities cannot be quoted at this time.

Likely container type:	Container	Waste packaged (%vol)	Waste loading (m ³)	Payload (m ³)	Number of packages
	Sellafield enhanced 3m ³ box	100.0	0.904	2.15	670

Likely container type comment: 1 skip of retrieved waste (max 1,400 litres) in = 1 package out.

Range in container waste volume: There will be considerable variability in unconditioned waste volume per package due to variations in skip loading and content. The actual number of packages produced is identified in the Sellafield Decommissioning Product and Secondary Waste Plan. 670 Boxes = 630 Boxes plus 40 from residuals

Other information on containers: Stainless Steel.

Likely conditioning matrix: BFS/OPC;PFA/OPC

Other information: -

Conditioned density (t/m³): ~2.0

Conditioned density comment: Density of conditioned waste will be fairly uniform.

Other information on conditioning: Waste matrix (as retrieved) will be in-filled with grout. A second pour of capping grout will be added. Void spaces between Skip wall and Box wall will be filled with grout.

Opportunities for alternative disposal routing: No

WASTE STREAM	2D08.1	MSSS Compartment 1
---------------------	---------------	---------------------------

Baseline Management Route	Opportunity Management Route	Stream volume (%)	Estimated Date that Opportunity will be realised	Opportunity Confidence	Comment
-	-	-	-	-	-

RADIOACTIVITY

Source:	Irradiated fuel, fuel components, reactor components and miscellaneous items contaminated with fuel residues.
Uncertainty:	Accuracy is generally good for fuel based residues, based on decanning records arisings and fuel carry over evaluations. For miscellaneous waste items, where records are incomplete, the accuracy is worse.
Definition of total alpha and total beta/gamma:	FISPIN generates data for approximately 2500 radionuclides. Only a proportion of these are considered in the UK inventory for this waste stream.
Measurement of radioactivities:	Isotopic specific activity (TBq/m ³) is derived by dividing the estimated total activity of the isotope by the total volume of the waste.
Other information:	Short-lived daughters are included in the "other" beta/gamma activity.

WASTE STREAM 2D08.1 MSSS Compartment 1

Nuclide	Mean radioactivity, TBq/m ³				Nuclide	Mean radioactivity, TBq/m ³			
	Waste at 1.4.2022	Bands and Code	Future arisings	Bands and Code		Waste at 1.4.2022	Bands and Code	Future arisings	Bands and Code
H 3	5.12E-02	AB 2			Gd 153	7.15E-28	AA 2		
Be 10	2.06E-07	AA 2			Ho 163	3.83E-12	AA 2		
C 14	4.68E-02	AB 2			Ho 166m	1.55E-08	AA 2		
Na 22					Tm 170		AA 2		
Al 26					Tm 171	1.07E-13	AA 2		
Cl 36	1.86E-06	AB 2			Lu 174				
Ar 39	4.10E-04	AB 2			Lu 176				
Ar 42					Hf 178n				
K 40	7.46E-09	AB 2			Hf 182	7.69E-17	AA 2		
Ca 41	5.69E-05	AB 2			Pt 193				
Mn 53	1.14E-09	AB 2			Tl 204	3.53E-06	AA 2		
Mn 54					Pb 205	5.92E-10	AA 2		
Fe 55	5.32E-04	AA 2			Pb 210	8.38E-10	AA 2		
Co 60	1.92E-01	AA 2			Bi 208				
Ni 59	6.44E-01	AA 2			Bi 210m	4.06E-14	AA 2		
Ni 63	5.55E+01	AA 2			Po 210	8.17E-10	AA 2		
Zn 65					Ra 223	2.04E-08	AA 2		
Se 79	4.83E-06	AA 2			Ra 225	1.15E-11	AA 2		
Kr 81					Ra 226	2.09E-09	AA 2		
Kr 85	2.63E-02	AA 2			Ra 228				
Rb 87	6.92E-10	AA 2			Ac 227	8.04E-09	AA 2		
Sr 90	1.82E+00	AA 2			Th 227	7.90E-09	AA 2		
Zr 93	1.55E-04	AA 2			Th 228	7.47E-08	AA 2		
Nb 91					Th 229	1.15E-11	AA 2		
Nb 92	4.47E-16	AA 2			Th 230	1.72E-07	AA 2		
Nb 93m	6.19E-03	AA 2			Th 232	4.03E-13	AA 2		
Nb 94	2.34E-03	AA 2			Th 234	3.53E-04	AA 2		
Mo 93	6.88E-03	AA 2			Pa 231	1.46E-08	AA 2		
Tc 97					Pa 233	8.24E-06	AA 2		
Tc 99	1.35E-03	AB 2			U 232	7.22E-08	AA 2		
Ru 106	8.36E-15	AA 2			U 233	2.97E-09	AA 2		
Pd 107	6.50E-06	AA 2			U 234	3.30E-04	AA 2		
Ag 108m	9.41E-10	AA 2			U 235	1.17E-05	AA 2		
Ag 110m	6.39E-25	AA 2			U 236	2.28E-05	AA 2		
Cd 109	2.15E-19	AA 2			U 238	3.53E-04	AA 2		
Cd 113m	2.05E-04	AA 2			Np 237	8.24E-06	AA 2		
Sn 119m					Pu 236	6.06E-12	AA 2		
Sn 121m	1.47E-05	AA 2			Pu 238	8.87E-03	AA 2		
Sn 123					Pu 239	1.13E-01	AA 2		
Sn 126	5.27E-05	AA 2			Pu 240	7.15E-02	AA 2		
Sb 125	1.19E-06	AA 2			Pu 241	3.98E-01	AA 2		
Sb 126	7.33E-06	AA 2			Pu 242	1.74E-05	AA 2		
Te 125m	2.91E-07	AA 2			Am 241	1.61E-01	AA 2		
Te 127m					Am 242m	9.95E-05	AA 2		
I 129	2.47E-06	AB 2			Am 243	2.32E-05	AA 2		
Cs 134	2.94E-08	AA 2			Cm 242	8.26E-05	AA 2		
Cs 135	6.42E-05	AA 2			Cm 243	6.16E-06	AA 2		
Cs 137	2.45E+00	AB 2			Cm 244	8.03E-05	AA 2		
Ba 133	1.60E-11	AA 2			Cm 245	2.57E-08	AA 2		
La 137					Cm 246	3.42E-09	AA 2		
La 138	5.28E-15	AA 2			Cm 248				
Ce 144	1.02E-18	AA 2			Cf 249	2.84E-14	AA 2		
Pm 145					Cf 250	9.89E-15	AA 2		
Pm 147	1.77E-05	AA 2			Cf 251	7.18E-16	AA 2		
Sm 147					Cf 252	6.39E-20	AA 2		
Sm 151	3.89E-02	AA 2			Other a	1.13E-01			
Eu 152	6.10E-05	AA 2			Other b/g	4.13E+00			
Eu 154	1.39E-03	AA 2			Total a	4.69E-01	AB 2	0	
Eu 155	1.08E-04	AA 2			Total b/g	6.53E+01	AB 2	0	

Bands (Upper and Lower)

- A a factor of 1.5
- B a factor of 3
- C a factor of 10
- D a factor of 100
- E a factor of 1000

Note: Bands quantify uncertainty in mean radioactivity.

Code

- 1 Measured activity
- 2 Derived activity (best estimate)
- 3 Derived activity (upper limit)
- 4 Not present
- 5 Present but not significant
- 6 Likely to be present but not assessed
- 7 Present in significant quantities but not determined
- 8 Not expected to be present in significant quantity