SITE Sellafield

SITE OWNER Nuclear Decommissioning Authority

Nο

WASTE CUSTODIAN Sellafield Limited

WASTE TYPE ILW

Is the waste subject to

WASTE VOLUMES

Scottish Policy:

Total future arisings: 0 m³

Total waste volume: 537.0 m³

Comment on volumes: There are no future arisings to this stream. Waste volume includes swarf/sludge,

miscellaneous beta/gamma waste, aggregate and associated water.

Uncertainty factors on Stock (upper): x 1.5 Arisings (upper) x

volumes: Stock (lower): x 0.66 Arisings (lower) x

WASTE SOURCE Swarf from Magnox decanning with small amounts of miscellaneous beta/gamma, mainly

from Windscale, some from other sites.

PHYSICAL CHARACTERISTICS

General description: Magnox cladding, much of which has been converted to a magnesium hydroxide sludge.

Some miscellaneous beta/gamma wastes (MBGW) are also present: these include tins/cans, filters, tools, pipework, pumps, valves, plastic items. The waste also includes some uranium most of which has corroded to oxides of uranium. A small volume of limestone aggregate is present at the bottom of the compartment. Swarf from Magnox decanning has reacted with the storage cover water to form corrosion products. Some

degradation of miscellaneous components waste will have occurred.

Physical components (%wt): Magnox swarf (8.61%), Magnox sludge (38.16%), Al/Al(OH)3 (0.01%), U/UO2/U3O8

(4.98%), MBGW (scrap, filters, cans, etc.) (1.68%), aggregate (limestone) (1.42%), water

(44.61%), others (0.53%). Includes interstitial liquor and cover liquor.

Sealed sources: Not yet determined.

Bulk density (t/m³): ~1.51

Comment on density: Bulk density is solids only - includes water content based on assumptions of draining

characteristics. Bulk density including interstitial liquor = $1.52t/m^3$, bulk density with cover liquor and interstitial liquor = $1.46t/m^3$. Average decanning debris is $\sim 1.6t/m^3$ and average

MBGW is ~3.2t/m3.

CHEMICAL COMPOSITION

General description and components (%wt):

Magnox swarf (8.61%), Magnox sludge (38.16%), Al/Al(OH)3 (0.01%), U/UO2/U3O8 (4.98%), MBGW (scrap, filters, cans, etc.) (1.68%), aggregate (limestone) (1.42%), water

(44.61%), others (0.53%). Includes interstitial liquor and cover liquor.

Chemical state: Neutral

Chemical form of H-3: Present in elemental and reacted forms. radionuclides: C-14: Present in elemental and reacted forms.

CI-36: Present in elemental and reacted forms. Se-79: Present in elemental and reacted forms. Tc-99: Present in elemental and reacted forms. I-129: Present in elemental and reacted forms.

Ra: Ra isotopes are present in less than trace amounts in fuel.

Th: Present in elemental and reacted forms.

U: Present in metallic and reacted forms (oxides and possibly hydride).

Np: Present in elemental and reacted forms. Pu: Present in metallic and mixed oxide forms.

Metals and alloys (%wt): No significant quantities of sheet metal, although small quantities of fabricated items are

present, e.g. paint tins, HEPA filters. Small pieces of Magnox metal present.

	(%wt)	Type(s) / Grade(s) with proportions	% of total C14
Stainless steel	~1.1		activity
Other ferrous metals	~3.6		
Iron	Р		
Aluminium	~0.01		
Beryllium	NE		
Cobalt	Р		
Copper	TR		
Lead	Р		
Magnox/Magnesium	~8.3	Predominantly Magnox.	
Nickel	TR		
Titanium	NE		
Uranium	~2.0		
Zinc	TR		
Zircaloy/Zirconium	~0.04		
Other metals	~0.05		
Organics (%wt): The organic content	of the was	ete is low.	
	(%wt)	Type(s) and comment	% of total C14 activity
Total cellulosics	~0.13		activity
Paper, cotton			
Wood			
Halogenated plastics	~0.07		
Total non-halogenated plastics	~0.02		
Condensation polymers			
Others			
Organic ion exchange materials	NE		
Total rubber	NE		
Halogenated rubber			
Non-halogenated rubber			
Hydrocarbons			
Oil or grease	<0.01		
Fuel	NE		
Asphalt/Tarmac (cont.coal tar)	NE		
Asphalt/Tarmac (no coal tar)	NE		
Bitumen	NE		
Others	NE		
Other organics	<0.01		
Other materials (%wt):			

	(%wt)	Type(s) and comment % of total C14 activity
Inorganic ion exchange materials	NE	activity
Inorganic sludges and flocs	~39.6	
Soil	NE	
Brick/Stone/Rubble	~1.4	
Cementitious material	NE	
Sand	NE	
Glass/Ceramics	~0.38	
Graphite	~0.02	
Desiccants/Catalysts	NE	
Asbestos	NE	
Non/low friable		
Moderately friable		
Highly friable		
Free aqueous liquids	~43.0	Includes interstitial liquor and cover liquor. Total waste volume 603 m3 with cover liquor.
Free non-aqueous liquids	NE	·
Powder/Ash	NE	
Inorganic anions (%wt): Hydroxides and car	bonates ar	re present.
	(%wt)	Type(s) and comment
-		7, - (-),
Fluoride	TR	
Chloride	TR	Alkali metal and alkaline earth chlorides.
lodide	TR	
Cyanide	NE 0.04	
Carbonate	~0.04	
Nitrate	TR	
Nitrite	TR	
Phosphate	TR	
Sulphate	TR	
Sulphide	TR	
Materials of interest for Waste contains ura waste acceptance criteria: as putrescible wast		nox and traces of uranium hydride. Cellulosics are present
	(%wt)	Type(s) and comment
Combustible metals	Р	
Low flash point liquids	0	
Explosive materials	0	
Phosphorus	0	
Hydrides	Р	Trace, passivated.
Biological etc. materials	0	
Biodegradable materials	Р	
Putrescible wastes	Р	Celluosics.

Non-putrescible wastes	Р	Organics - plastics and rubbers.
Corrosive materials	0	
Pyrophoric materials	0	
Generating toxic gases	0	
Reacting with water	Р	<5%.
Higher activity particles	Р	Radioactive sludges and solids.
Soluble solids as bulk chemical compounds	Р	Elemental and hydroxides, oxides and carbonates.
Hazardous substances / - non hazardous pollutants:		
	(%wt)	Type(s) and comment
Acrylamide		
Benzene	0	
Chlorinated solvents		
Formaldehyde		
Organometallics		
Phenol	0	
Styrene		
Tri-butyl phosphate	0	
Other organophosphates		
Vinyl chloride	0	
Arsenic	0	
Barium		
Boron	NE	Traces - elemental and borates.
Boron (in Boral)		
Boron (non-Boral)		
Cadmium	0	
Caesium		
Selenium	0	
Chromium	Р	In steels.
Molybdenum	Р	In steels.
Thallium		
Tin	Р	Traces.
Vanadium	Р	In steels.
Mercury compounds		
Others	NE	
Electronic Electrical Equipment (EEE)		
EEE Type 1		
EEE Type 2		
EEE Type 3		
EEE Type 4		
EEE Type 5		

0 1 :	1 (0/ 1)	N					
Complexing age	nts (%wt):	Not yet determined					
		(%	6wt) Ty	/pe(s) and c	omment		
EC)TA						
DF	PTA						
NT	A						
Ро	lycarboxylic ad	cids					
Oti	her organic co	mplexants					
То	tal complexing	agents					
Potential for the contain discrete i		Not yet determined.					
PACKAGING A	ND CONDIT	IONING					
Conditioning method: The waste will be retrieved from MSSS and stored in an urperiod of 70 years. Following the period of storage it will be and conditioned for ultimate disposal to the GDF.							
Plant Name:		TBD					
Location:		Sellafield					
Plant startup date	э:	2070-2090					
Total capacity (m³/y incoming w	aste):	-					
Target start date for - packaging this stream:							
Throughput for th (m³/y incoming w		-					
Other information	n:	The plant has not yet be quoted at this time.	en design	ed. Hence, 1	throughpu	ts & capacit	ies cannot be
Likely container type:	Container		Wast packag (%vo	ed load	/aste ling (m³)	Payload (m³)	Number of packages
	Sellafield enhanced 3m³ box		100.	0 0).658	2.15	816
Likely container t comment:	ype	1 skip of retrieved waste	(max 1,40	00 litres) in =	= 1 packaç	ge out.	
Range in container waste volume:		There will be considerable variability in unconditioned waste volume per package due to variations in skip loading and content. The actual number of packages produced is identified in the Sellafield Decommissioning Product and Secondary Waste Plan.816					

Boxes = 777 Boxes plus 39 from residuals.

Other information on containers:

Stainless Steel.

Likely conditioning matrix: BFS/OPC;PFA/OPC

Conditioned density (t/m³):

Other information:

~2.0

Conditioned density

comment:

Density of conditioned waste will be fairly uniform.

Other information on

conditioning:

Waste matrix (as retrieved) will be in-filled with grout. A second pour of capping grout will be added. Void spaces between Skip wall and Box wall will be filled with grout.

Opportunities for alternative

disposal routing:

No

Estimated

Date that Opportunity Opportunity Baseline Stream Comment Opportunity Management Route Management Route Confidence volume (%) will be realised

RADIOACTIVITY

Sr-90, Cs-137, Co-60, Pu-239, Pu-240, Pu-241, Am-241. Source:

Uncertainty: Accuracy is generally good - based on decanning records arisings and fuel carry over

evaluations.

Definition of total alpha

and total beta/gamma:

FISPIN generates data for approximately 2500 radionuclides. Only a proportion of these are considered in the UK inventory for this waste stream.

Measurement of Isotopic specific activity (TBq/m³) is derived by dividing the estimated total activity of the

radioactivities: isotope by the total volume of the waste.

Other information: Short-lived daughters are included in the "other" beta/gamma activity.

	Mean radioactivity, TBq/m³			Mean radioactivity, TBq/m³					
	Waste at	Bands and	Future	Bands and		Waste at	Bands and	Future	Bands and
Nuclide	1.4.2022	Code	arisings	Code	Nuclide	1.4.2022	Code	arisings	Code
H 3	2.64E-02	AB 2			Gd 153	6.04E-24	AA 2		
Be 10	3.65E-07	AA 2			Ho 163	2.39E-11	AA 2		
C 14	8.33E-03	AB 2			Ho 166m	1.78E-07	AA 2		
Na 22					Tm 170	2.00E-44	AA 2		
Al 26	0.045.07	4.5. 0			Tm 171	1.08E-11	AA 2		
CI 36	2.84E-07	AB 2			Lu 174				
Ar 39 Ar 42	5.69E-05	AB 2			Lu 176 Hf 178n				
K 40	1.43E-08	AB 2			Hf 182	4.65E-16	AA 2		
Ca 41	1.52E-04	AB 2			Pt 193	4.03L-10	AA 2		
Mn 53	1.86E-10	AB 2			TI 204	2.88E-06	AA 2		
Mn 54	1.002 10	712 2			Pb 205	4.11E-11	AA 2		
Fe 55	1.04E-03	AA 2			Pb 210	1.50E-09	AA 2		
Co 60	9.83E-02	AA 2			Bi 208				
Ni 59	8.44E-02	AA 2			Bi 210m	2.82E-15	AA 2		
Ni 63	7.82E+00	AA 2			Po 210	1.44E-09	AA 2		
Zn 65					Ra 223	4.25E-08	AA 2		
Se 79	2.19E-05	AA 2			Ra 225	4.79E-11	AA 2		
Kr 81					Ra 226	4.07E-09	AA 2		
Kr 85	1.92E-01	AA 2			Ra 228				
Rb 87	8.65E-11	AA 2			Ac 227	1.67E-08	AA 2		
Sr 90	9.31E+00	AA 2			Th 227	1.65E-08	AA 2		
Zr 93	6.86E-04	AA 2			Th 228	6.37E-07	AA 2		
Nb 91	0.705.45				Th 229	4.82E-11	AA 2		
Nb 92	2.78E-15	AA 2			Th 230	3.82E-07	AA 2		
Nb 93m Nb 94	1.19E-03 4.49E-04	AA 2 AA 2			Th 232 Th 234	1.99E-12 9.51E-04	AA 2 AA 2		
Mo 93	7.23E-04	AA 2			Pa 231	3.10E-08	AA 2		
Tc 97	7.202 01	707 2			Pa 233	4.34E-05	AA 2		
Tc 99	5.53E-03	AB 2			U 232	6.21E-07	AA 2		
Ru 106	4.82E-12	AA 2			U 233	1.33E-08	AA 2		
Pd 107	3.87E-05	AA 2			U 234	8.50E-04	AA 2		
Ag 108m	5.91E-09	AA 2			U 235	2.50E-05	AA 2		
Ag 110m	3.86E-21	AA 2			U 236	9.09E-05	AA 2		
Cd 109	3.13E-19	AA 2			U 238	9.51E-04	AA 2		
Cd 113m	5.82E-05	AA 2			Np 237	4.34E-05	AA 2		
Sn 119m					Pu 236	1.76E-10	AA 2		
Sn 121m	2.40E-06	AA 2			Pu 238	1.02E-01	AA 2		
Sn 123	2.87E-04	A A 2			Pu 239	3.99E-01	AA 2		
Sn 126 Sb 125	4.68E-05	AA 2 AA 2			Pu 240 Pu 241	3.97E-01 4.27E+00	AA 2 AA 2		
Sb 126	4.03E-05	AA 2			Pu 242	2.05E-04	AA 2		
Te 125m	1.15E-05	AA 2			Am 241	1.12E+00	AA 2		
Te 127m		=			Am 242m	1.43E-03	AA 2		
l 129	1.21E-05	AB 2			Am 243	3.98E-04	AA 2		
Cs 134	2.93E-06	AA 2			Cm 242	1.19E-03	AA 2		
Cs 135	3.02E-04	AA 2			Cm 243	1.51E-04	AA 2		
Cs 137	1.36E+01	AB 2			Cm 244	1.81E-03	AA 2		
Ba 133	1.68E-10	AA 2			Cm 245	2.79E-07	AA 2		
La 137	0.405				Cm 246	2.73E-08	AA 2		
La 138	3.43E-14	AA 2			Cm 248	1 555 40	^ ^ ^		
Ce 144	1.18E-15	AA 2			Cf 249	1.55E-13	AA 2		
Pm 145 Pm 147	5.79E-04	AA 2			Cf 250 Cf 251	8.88E-14 3.80E-15	AA 2 AA 2		
Sm 147	5.7 JL-04	111 Z			Cf 251	4.60E-18	AA 2		
Sm 151	1.37E-01	AA 2			Other a	4.01E-01	=		
Eu 152	5.33E-04	AA 2			Other b/g	2.23E+01			
Eu 154	2.28E-02	AA 2			Total a	2.42E+00	AB 2	0	
Eu 155	1.48E-03	AA 2			Total b/g	5.79E+01	AB 2	0	
	1		ı			ı		I	

Bands (Upper and Lower)

A a factor of 1.5 B a factor of 3 C a factor of 10 D a factor of 100

D a factor of 100 E a factor of 1000

Note: Bands quantify uncertainty in mean radioactivity.

Code

- 1 Measured activity
- 2 Derived activity (best estimate)
- 3 Derived activity (upper limit)
- 4 Not present
- 5 Present but not significant
- 6 Likely to be present but not assessed
- 7 Present in significant quantities but not determined