SITE Sellafield

SITE OWNER **Nuclear Decommissioning Authority**

WASTE CUSTODIAN Sellafield Limited

LLW **WASTE TYPE**

Is the waste subject to

Scottish Policy:

Nο

WASTE VOLUMES

| | | Reported |
|------------------------|----------------------|--------------------|
| Stocks: | At 1.4.2022 | 0 m³ |
| Future arisings - | 1.4.2022 - 31.3.2023 | 8.3 m ³ |
| | 1.4.2023 - 31.3.2024 | 8.3 m ³ |
| | 1.4.2024 - 31.3.2025 | 8.3 m ³ |
| | 1.4.2025 - 31.3.2026 | 8.3 m ³ |
| | 1.4.2026 - 31.3.2027 | 8.3 m ³ |
| | 1.4.2027 - 31.3.2028 | 8.3 m ³ |
| | 1.4.2028 - 31.3.2029 | 8.3 m ³ |
| | 1.4.2029 - 31.3.2030 | 8.3 m ³ |
| | 1.4.2030 - 31.3.2031 | 8.3 m ³ |
| | 1.4.2031 - 31.3.2032 | 8.3 m ³ |
| | 1.4.2032 - 31.3.2033 | 8.3 m ³ |
| | 1.4.2033 - 31.3.2034 | 8.3 m ³ |
| | 1.4.2034 - 31.3.2035 | 8.3 m ³ |
| | 1.4.2035 - 31.3.2036 | 8.3 m ³ |
| Total future arisings: | | 116.5 m³ |
| Total waste volume: | | 116.5 m³ |

Arisings are sourced from REM_TP_0116A and are based on the latest five-year forecasts Comment on volumes:

from the Waste Forecasting database. The overall timescale for waste arising are informed

by the Sellafield Site Master Timeline. Uncertainty information is notional.

Uncertainty factors on Stock (upper): Arisings (upper) volumes: Stock (lower): Arisings (lower) x 0.5

WASTE SOURCE Waste arises as a result of routine operations and maintenance in SETP.

PHYSICAL CHARACTERISTICS

The waste will mostly be a mixture of metallic wastes associated with maintenance and General description:

redundant plant items as well as secondary wastes associated with operatons and maintenance. The waste has not undergone any changes since it was generated.

Metals (43.2%), Concrete/Rubble (5%), Soil (1%), Wood (2%), Rubber (2%), Halogenated Physical components (%wt):

Plastics (17.9%), Non-Halogenated Plastics (17.9%), Hydrocarbons (2.2%), Other

Organics (8.2%), Asbestos (0.4%) and Other (0.2%).

Sealed sources: The waste does not contain sealed sources.

0.633 Bulk density (t/m³):

Comment on density: The total bulk density is derived from REM_TP_0116A and is based on the five-year

forecast from the Waste Forecasting Database.

CHEMICAL COMPOSITION

General description and

components (%wt):

Metals (43.2%), Concrete/Rubble (5%), Soil (1%), Wood (2%), Rubber (2%), Halogenated

Plastics (17.9%), Non-Halogenated Plastics (17.9%), Hydrocarbons (2.2%), Other

Organics (8.2%), Asbestos (0.4%) and Other (0.2%).

Chemical state: Neutral

Chemical form of radionuclides:

Metals and alloys (%wt): Metal thickness not specified.

| | (%wt) | Type(s) / Grade(s) with proportions | % of total C14 |
|--------------------------------|-------|-------------------------------------|-------------------------|
| Stainless steel | 6.3 | | activity |
| Other ferrous metals | 23.2 | | |
| Iron | 7.1 | | |
| Aluminium | 0.18 | | |
| Beryllium | 0 | | |
| Cobalt | 0 | | |
| Copper | 0.18 | | |
| Lead | 6.3 | | |
| Magnox/Magnesium | 0 | | |
| Nickel | 0 | | |
| Titanium | 0 | | |
| Uranium | 0 | | |
| Zinc | 0 | | |
| Zircaloy/Zirconium | 0 | | |
| Other metals | 0 | | |
| Organics (%wt): | | | |
| | (%wt) | Type(s) and comment | % of total C14 activity |
| Total cellulosics | 2.0 | | activity |
| Paper, cotton | 0 | | |
| Wood | 2.0 | | |
| Halogenated plastics | 17.9 | | |
| Total non-halogenated plastics | 17.9 | | |
| Condensation polymers | 0 | | |
| Others | 0 | | |
| Organic ion exchange materials | 0 | | |
| Total rubber | 2.0 | | |
| Halogenated rubber | 0 | | |
| Non-halogenated rubber | 0 | | |
| Hydrocarbons | 2.2 | | |
| Oil or grease | 0 | | |
| Fuel | 0 | | |
| Asphalt/Tarmac (cont.coal tar) | 1.8 | | |
| Asphalt/Tarmac (no coal tar) | 0.09 | | |
| Bitumen | 0.36 | | |
| Others | 0 | | |
| Other organics | 8.2 | | |
| Other materials (%wt): | | | |

| | (%wt) | Type(s) and comment | % of total C14 activity |
|--|-------|---------------------|-------------------------|
| Inorganic ion exchange materials | 0 | | |
| Inorganic sludges and flocs | 0 | | |
| Soil | 1.0 | | |
| Brick/Stone/Rubble | 5.0 | | |
| Cementitious material | 0 | | |
| Sand | 0 | | |
| Glass/Ceramics | 0.18 | | |
| Graphite | 0 | | |
| Desiccants/Catalysts | 0 | | |
| Asbestos | 0.45 | | |
| Non/low friable | 0.18 | | |
| Moderately friable | 0.18 | | |
| Highly friable | 0.09 | | |
| Free aqueous liquids | 0 | | |
| Free non-aqueous liquids | 0 | | |
| Powder/Ash | 0 | | |
| Inorganic anions (%wt): | | | |
| | (%wt) | Type(s) and comment | |
| Fluoride | 0 | | |
| Chloride | 0 | | |
| lodide | 0 | | |
| Cyanide | 0 | | |
| Carbonate | 0 | | |
| Nitrate | 0 | | |
| Nitrite | 0 | | |
| Phosphate | 0 | | |
| Sulphate | 0 | | |
| Sulphide | 0 | | |
| Materials of interest for - waste acceptance criteria: | | | |
| | (%wt) | Type(s) and comment | |
| Combustible metals | 0 | | |
| Low flash point liquids | 0 | | |
| Explosive materials | 0 | | |
| Phosphorus | 0 | | |
| Hydrides | 0 | | |
| Biological etc. materials | 0 | | |
| Biodegradable materials | 4.0 | | |
| Putrescible wastes | 2.0 | | |
| Non-putrescible wastes | 2.0 | | |

| | Corrosive materials | 0 | |
|-------------|---|-------|---------------------------------|
| | Pyrophoric materials | 0 | |
| | Generating toxic gases | 0 | |
| | Reacting with water | 0.18 | |
| | Higher activity particles | 0 | |
| | Soluble solids as bulk chemical compounds | 0 | |
| Hazardous s | | | |
| | | (%wt) | Type(s) and comment |
| | Acrylamide | 0 | |
| | Benzene | 0 | |
| | Chlorinated solvents | 0 | |
| | Formaldehyde | 0 | |
| | Organometallics | 0 | |
| | Phenol | 0 | |
| | Styrene | 0 | |
| | Tri-butyl phosphate | 0 | |
| | Other organophosphates | 0 | |
| | Vinyl chloride | 0 | |
| | Arsenic | 0 | |
| | Barium | 0 | |
| | Boron | 0 | |
| | Boron (in Boral) | 0 | |
| | Boron (non-Boral) | 0 | |
| | Cadmium | 0 | |
| | Caesium | 0 | |
| | Selenium | 0 | |
| | Chromium | 0 | |
| | Molybdenum | 0 | |
| | Thallium | 0 | |
| | Tin | 0 | |
| | Vanadium | 0.13 | |
| | Mercury compounds | 0 | |
| | Others | 0 | |
| | Electronic Electrical Equipment (EEE) | | |
| | EEE Type 1 | | 200 Items over the next 5 years |
| | EEE Type 2 | | 200 Items over the next 5 years |
| | EEE Type 3 | | 100 Items over the next 5 years |
| | EEE Type 4 | | 100 Items over the next 5 years |
| | EEE Type 5 | | 50 Items over the next 5 years |
| | | | |

Complexing agents (%wt): Yes

Potential for the waste to contain discrete items:

Yes. Pumps, motors and hand tools.

TREATMENT, PACKAGING AND DISPOSAL

Planned on-site / off-site treatment(s):

| Treatment | On-site / Off site | Stream volume % |
|-----------------------|-----------------------|-----------------|
| Low force compaction | | |
| Supercompaction (HFC) | On-site | 14.4 |
| Incineration | Off-site | 40.9 |
| Solidification | | |
| Decontamination | | |
| Metal treatment | Off-site | 11.1 |
| Size reduction | | |
| Decay storage | | |
| Recyling / reuse | | |
| Other / various | | |
| None | | 33.6 |

Comment on planned treatments:

All high force compaction takes place in WAMAC. For Inventory purposes, it is assumed that supercompaction will continue after the closure of WAMAC in 2028 $\,$

Disposal Routes:

| Disposal Route | Stream volume % | Disposal density t/m3 |
|--|-----------------|--------------------------|
| Expected to be consigned to the LLW Repository | 48.0 | 0.88 |
| Expected to be consigned to a Landfill Facility | | |
| Expected to be consigned to an On-Site Disposal Facility | | |
| Expected to be consigned to an Incineration Facility | 40.9 | 0.14 |
| Expected to be consigned to a Metal Treatment Facility | 11.1 | 1.4 |
| Expected to be consigned as Out of Scope | | |
| Expected to be recycled / reused | | |
| Disposal route not known | | |

Classification codes for waste expected to be consigned to a landfill facility:

Upcoming (2022/23-2024/25) Waste Routing (if expected to change from above):

| Disposal Route | Stream volume % | | | |
|--|-----------------|---------|---------|--|
| Disposal Notice | 2022/23 | 2023/24 | 2024/25 | |
| Expected to be consigned to the LLW Repository Expected to be consigned to a Landfill Facility Expected to be consigned to an On-Site Disposal Facility Expected to be consigned to an Incineration Facility Expected to be consigned to a Metal Treatment Facility Expected to be consigned as Out of Scope Expected to be recycled / reused Disposal route not known | | | | |

Opportunities for alternative disposal routing: No

Baseline Opportunity Stream Date that Opportunity Confidence Wanagement Route Management Route Volume (%)

Estimated Opportunity Confidence will be realised

Waste Packaging for Disposal:

| Container | Stream volume % | Waste loading m ³ | Number of packages | |
|--|-----------------|------------------------------|--------------------|--|
| 1/3 Height IP-1 ISO 2/3 Height IP-2 ISO | | | | |
| 1/2 Height WAMAC IP-2 ISO | 14.4 | 59.28 | < 1 | |
| 1/2 Height IP-2 Disposal/Re-usable ISO | 33.6 | 10 | 4 | |
| 2m box (no shielding) | | | | |
| 4m box (no shielding) | | | | |
| Other | | | | |

Other information: -

Waste Planned for Disposal at the LLW Repository:

Container voidage: -

Waste Characterisation

Form (WCH):

The waste meets the LLWR's Waste Acceptance Criteria (WAC).

The waste has a current WCH.

Differences exist between Inventory information and current WCH.

Materials and radioactivity data has been taken from the current WCH, but data on waste volumes and waste routes is based on the Waste Forecasting database as

this information is more recent.

Waste consigned for disposal to LLWR in year of generation:

Yes.

Non-Containerised Waste for In-Vault Grouting: (Not applicable to this waste stream)

Stream volume (%):

Waste stream variation: -

Bounding cuboidal volume:

Inaccessible voidage: -

Other information: -

RADIOACTIVITY

Source: The activity originates from effluents that in turn arise from a variety of site operations.

Uncertainty: The uncertainty associated with the fingerprinting analysis is likely to be low, however the

volumes and total activity information (and possibly some other assumptions) are likely to

be more notional and thus more uncertain.

Definition of total alpha and total beta/gamma:

Where totals are shown on the table of radionuclide activities they are the sums of the listed alpha or beta/gamma emitting radionuclides plus 'other alpha' or 'other beta/gamma'.

Measurement of radioactivities:

Specific activity data is based on data in the corresponding WCH, which in turn maps an

estimated total activity to an analytically derived radionuclide fingerprint.

Other information: The radionuclides have been taken from REM_TP_0116A and are based on the current

WCH (LLWR Ref: 1S-1S-0-WCH-0-3695 version 7).

| | Mean radioactivity, TBq/m³ | | | | Mean radioactivity, TBq/m³ | | | | |
|-------------------|----------------------------|-----------|----------------------|--------------|----------------------------|----------|-----------|----------|--------------|
| Nuclide | Waste at | Bands and | Future | Bands and | Nuclide | Waste at | Bands and | Future | Bands and |
| H 3 | 1.4.2022 | Code | arisings 3.16E-06 | Code CC 2 | Gd 153 | 1.4.2022 | Code | arisings | Code |
| Be 10 | | | 3.10E-00 | CC 2 | Ho 163 | | | | |
| C 14 | | | 2.28E-06 | CC 2 | Ho 166m | | | | |
| Na 22 | | | 2.202 00 | 00 2 | Tm 170 | | | | |
| Al 26 | | | | | Tm 171 | | | | |
| CI 36 | | | | | Lu 174 | | | | |
| Ar 39 | | | | | Lu 176 | | | | |
| Ar 42 | | | | | Hf 178n | | | | |
| K 40 | | | | | Hf 182 | | | | |
| Ca 41 | | | | | Pt 193 | | | | |
| Mn 53 | | | | | TI 204 | | | | |
| Mn 54 | | | | | Pb 205 | | | | |
| Fe 55 | | | | | Pb 210 | | | | |
| Co 60 | | | 5.43E-08 | CC 2 | Bi 208 | | | | |
| Ni 59 | | | | | Bi 210m | | | | |
| Ni 63 | | | | | Po 210 | | | | |
| Zn 65 | | | | | Ra 223 | | | | |
| Se 79 | | | | | Ra 225 | | | | |
| Kr 81 | | | | | Ra 226 | | | | |
| Kr 85 | | | | | Ra 228 | | | | |
| Rb 87 | | | | | Ac 227 | | | | |
| Sr 90 | | | 1.21E-05 | CC 2 | Th 227 | | | | |
| Zr 93 | | | | | Th 228 | | | | |
| Nb 91 | | | | | Th 229 | | | | |
| Nb 92 | | | | | Th 230 | | | | |
| Nb 93m | | | | | Th 232 | | | | |
| Nb 94 | | | | | Th 234 | | | | |
| Mo 93 | | | | | Pa 231 | | | | |
| Tc 97 | | | | | Pa 233 | | | | |
| Tc 99 | | | 3.73E-08 | CC 2 | U 232 | | | | |
| Ru 106 | | | 1.59E-07 | CC 2 | U 233 | | | 0.075.00 | 00.0 |
| Pd 107 | | | | | U 234 | | | 2.37E-08 | CC 2 |
| Ag 108m | | | | | U 235 | | | 9.04E-10 | CC 2 |
| Ag 110m | | | | | U 236 U 238 | | | 3.39E-09 | CC 2 CC 2 |
| Cd 109 Cd 113m | | | | | Np 237 | | | 2.71E-08 | 00 2 |
| Sn 119m | | | | | Pu 236 | | | | |
| Sn 121m | | | | | Pu 238 | | | 6.44E-08 | CC 2 |
| Sn 123 | | | | | Pu 239 | | | 6.78E-08 | CC 2 |
| Sn 126 | | | | | Pu 240 | | | 6.78E-08 | CC 2 |
| Sb 125 | | | 1.29E-07 | CC 2 | Pu 241 | | | 3.68E-06 | CC 2 |
| Sb 126 | | | | 00 2 | Pu 242 | | | 0.002 00 | 00 2 |
| Te 125m | | | | | Am 241 | | | 1.02E-07 | CC 2 |
| Te 127m | | | | | Am 242m | | | | - |
| I 129 | | | | | Am 243 | | | | |
| Cs 134 | | | 7.80E-08 | CC 2 | Cm 242 | | | | |
| Cs 135 | | | | | Cm 243 | | | | |
| Cs 137 | | | 1.19E-05 | CC 2 | Cm 244 | | | 1.36E-08 | CC 2 |
| Ba 133 | | | | | Cm 245 | | | | |
| La 137 | | | | | Cm 246 | | | | |
| La 138 | | | | | Cm 248 | | | | |
| Ce 144 | | | | | Cf 249 | | | | |
| Pm 145 | | | | | Cf 250 | | | | |
| Pm 147 | | | | | Cf 251 | | | | |
| Sm 147 | | | | | Cf 252 | | | | |
| Sm 151 | | | | | Other a | | | | |
| Eu 152 | | | | | Other b/g | | | | |
| Eu 154 | | | | | Total a | 0 | | 3.71E-07 | CC 2 |
| Eu 155 | | | | | Total b/g | 0 | | 3.35E-05 | CC 2 |
| | • | | | | | | | • | |

Bands (Upper and Lower)

A a factor of 1.5 B a factor of 3 C a factor of 10 D a factor of 100

E a factor of 1000

Note: Bands quantify uncertainty in mean radioactivity.

Code

- 1 Measured activity
- 2 Derived activity (best estimate)
- 3 Derived activity (upper limit)
- 4 Not present
- 5 Present but not significant
- 6 Likely to be present but not assessed
- 7 Present in significant quantities but not determined