SITE Berkeley

SITE OWNER Nuclear Decommissioning Authority

WASTE CUSTODIAN Magnox Limited

WASTE TYPE ILW

Is the waste subject to Scottish Policy:

No

WASTE VOLUMES

Reported

Stocks: At 1.4.2022...... 0.1 m³

Total future arisings: 0 m³

Total waste volume: 0.1 m³

Comment on volumes: There will be no further arisings of this waste stream. The volume given is the volume of

Magnox fuel element debris as held in drums.

Uncertainty factors on volumes:

Stock (upper): x 1.1 Stock (lower): x 0.9 Arisings (upper) x

Arisings (lower) x

WASTE SOURCE

The waste arose from the programme of post irradiation examination of Magnox fuel elements that was undertaken within the research facilities previously on the site.

PHYSICAL CHARACTERISTICS

General description: The waste is predominantly made up of Magnox metal generated from the post irradiation

examination work, although it could include other fuel element components. The largest arisings are believed to be splitter cages. There may be a significant quantity of Magnox fuel canning material contaminated by fuel in the debris. There will also be some Sintox discs, along with graphite, stainless steel and zirconium. The waste may be contaminated with Ternary Eutectic Chloride (TEC) (a fire extinguishant). All of the waste is containerised in mild steel black cans. The waste containers will be processed with this waste stream.

There are no large items that may require special handling.

Physical components (%wt): The vast majority of the FED from Berkeley Centre is Magnox (98%wt) together with

possible small quantities of zirconium (1.6%wt), Sintox discs (sintered alumina) (0.1%wt), other components (Nimonic springs, graphite, stainless steel) (0.23%wt) and <<0.1%wt thermocouples. There will also be small amounts of mild steel from the containers, and

there may be traces of Ternary Eutectic Chloride present.

Sealed sources: The waste does not contain sealed sources.

Bulk density (t/m³): 0.54

Comment on density: Density is calculated from the total mass of the loose waste, and the containerised waste

(including the containers), and the volume.

CHEMICAL COMPOSITION

General description and components (%wt):

The component breakdown of the FED from Berkeley Centre is 98%wt Magnox, 1.6%wt Zirconium, 0.1%wt Sintox discs (sintered alumina), 0.23%wt. Other components (Nimonic springs, graphite, stainless steel) and <<0.1%wt Thermocouples. There will also be small amounts of mild steel from the containers, and there may be traces of Ternary Eutectic

Chloride present.

Chemical state: Alkali

Chemical form of radionuclides:

H-3: Tritium is expected to be present as surface contamination, possibly as water, but

perhaps in the form of other inorganic or organic compounds.

C-14: Carbon 14 will probably be present as graphite.

CI-36: Chlorine 36 incorporated in the Magnox may be associated with barium impurity

(barium chloride).

Se-79: The selenium content is insignificant. Tc-99: The technetium content is insignificant. Ra: Radium isotope content is insignificant. Th: The thorium isotope content is insignificant.

U: Chemical form of U isotopes has not been determined but may be oxides.

Np: The neptunium content is insignificant.

Pu: Chemical form of plutonium isotopes has not been determined but may be plutonium

oxides.

Metals and alloys (%wt): Bulk items are not expected to be present. Nearly all of the waste is expected to be only 1 or 2 mm thick. Type(s) / Grade(s) with proportions (%wt) % of total C14 activity Stainless steel..... < 0.20 Other ferrous metals..... ΝE Mild steel will be present as the waste container. Iron. Aluminium..... Beryllium..... TR Cobalt..... Copper...... 0 Lead...... 0 0.8 wt% aluminium as an alloying constituent of Magnox. Nickel.....~0.20 Nimonic. Titanium..... Uranium..... Zinc..... Zircaloy/Zirconium..... Other metals..... "Other" metals could include thermocouple wires. Organics (%wt): Organics may be present in trace quantities. Organic ion exchange resin would be phenol formaldehyde based. (%wt) Type(s) and comment % of total C14 activity Total cellulosics..... 0 Paper, cotton..... 0 Wood..... 0 Halogenated plastics 0 Total non-halogenated plastics..... 0 Condensation polymers..... 0 Others..... 0 Organic ion exchange materials.... TR 0 Total rubber..... 0 Halogenated rubber Non-halogenated rubber..... 0 Hydrocarbons..... Oil or grease Fuel..... Asphalt/Tarmac (cont.coal tar)...

Other materials (%wt): Traces of graphite may be present.

Asphalt/Tarmac (no coal tar).....

Bitumen.....

Others....

Other organics....

TR

			(%wt)	Type(s) and comment	% of total C14 activity
	Inorganic ion ex	xchange materials	TR		•
	Inorganic sludg	es and flocs	0		
	Soil		0		
	Brick/Stone/Rul	bble	0		
	Cementitious m	naterial	0		
	Sand				
	Glass/Ceramics	3	0		
	Graphite		TR		
	Desiccants/Cat	alysts			
	Asbestos		0		
	Non/low fria	ble			
	Moderately	friable			
	Highly friabl	e			
	Free aqueous l	iquids	0		
	Free non-aqueo	ous liquids	0		
	Powder/Ash		Р		
Inorganic an	ions (%wt):	Inorganic anions a	e not expe	cted to be present at greater than	trace concentrations.
			(%wt)	Type(s) and comment	
	Fluoride		TR		
	Chloride		TR		
	lodide		0		
	Cyanide		0		
	Carbonate		TR		
	Nitrate		TR		
	Nitrite		TR		
	Phosphate		TR		
	Sulphate		TR		
	Sulphide		0		
Materials of i waste accep	interest for tance criteria:	Eutectic Chloride (ı	made up of e (25-51wt ^o	contaminated with the fire extingu Sodium chloride (20-35wt%, Pota %)). This may be of potential cond waste package.	assium chloride (29-40wt%)
			(%wt)	Type(s) and comment	
	Combustible me	etals	98.0		
	Low flash point liquids				
Explosive materials			0		
	Phosphorus		0		
	Hydrides		0		
	Biological etc. r	naterials	0		
	Biodegradable	materials	0		
	Putrescible w	vaetae	0		

Non-putrescible wastes		
Corrosive materials	0	
Pyrophoric materials	0	
Generating toxic gases	0	
Reacting with water	98.0	
Higher activity particles		
Soluble solids as bulk chemical		
compounds		
Hazardous substances / None expected. non hazardous pollutants:		
	(%wt)	Type(s) and comment
Acrylamide		
Benzene		
Chlorinated solvents		
Formaldehyde		
Organometallics		
Phenol		
Styrene		
Tri-butyl phosphate		
Other organophosphates		
Vinyl chloride		
Arsenic		
Barium		
Boron	0	
Boron (in Boral)		
Boron (non-Boral)		
Cadmium		
Caesium		
Selenium		
Chromium		
Molybdenum		
Thallium		
Tin		
Vanadium		
Mercury compounds		
Others		
Electronic Electrical Equipment (EEE	·)	
EEE Type 1		
EEE Type 2		
EEE Type 3		
EEE Type 4		
EEE Type 5		

Complexing ag	gents (%wt):	Yes							
		(%	6wt)	Type(s)	and comment				
E	DTA								
	PTA								
N	ITA								
P	olycarboxylic ac	ids							
C	Other organic cor	mplexants							
Т	otal complexing	agents TI	₹						
Potential for the contain discrete		Yes. In & of itself not a (see Nimonic/Others)	a DI; Wi	ll likely co	ontain "rogue" it	tems (HDRIs)	that will be		
PACKAGING	AND CONDIT	IONING							
Conditioning method:		This stream is to be co-packaged with 9A64, 9A83, 9A84, 9A33, 9A34, 9A35, 9A41, 9A42, 9A43, 9A49, 9A50, 9A51, 9A54, 9A55, 9A56, 9A74. Packages are assigned to 9A33/C, 9A34, 9A74.							
Plant Name:		-							
Location:		Berkeley Site							
Plant startup date: Total capacity		-							
•		-							
Target start dat packaging this		-							
Throughput for this stream (m³/y incoming waste):		-							
Other information:		-							
Likely contained type:	r Container		pack	aste aged svol)	Waste loading (m³)	Payload (m³)	Number of packages		
Likely container comment:	r type	-							
Range in container waste volume:		-							
Other information on containers:		-							
Likely conditioning matrix: Other information:		-							
Other information: Conditioned density (t/m³):		-							
Conditioned density comment:		-							
Other information on conditioning:		-							
Opportunities for alternative disposal routing:		-							

Baseline Opportunity Stream Date that Opportunity
Management Route Management Route volume (%)

Baseline Opportunity Opportunity Confidence will be realised

RADIOACTIVITY

Source: Activation when the associated fuel elements were irradiated, of nuclides incorporated in

the Magnox and other materials, contamination by fission products and actinides during

fuel element examination at Berkeley Technology Centre.

Uncertainty: The values quoted are indicative of the activities that might be expected.

Definition of total alpha and total beta/gamma:

Where totals are shown on the table of radionuclide activities they are the sums of the listed alpha or beta/gamma emitting radionuclides plus 'other alpha' or 'other beta/gamma'.

Measurement of radioactivities:

Values were derived from measurements, calculations of induced activity and estimates of

likely contamination.

Other information:

	ı	Mean radioac	tivity, TBq/m³				Mean radioactivity, TBq/m³			
Nuclide	Waste at 1.4.2022	Bands and Code	Future arisings	Bands and Code	Nuclide	Waste at 1.4.2022	Bands and Code	Future arisings	Bands and Code	
H 3	8.63E-03	CC 2			Gd 153		8			
Be 10	2E-07	CC 2			Ho 163		8			
C 14	9.99E-05	CC 2			Ho 166m		8			
Na 22		8			Tm 170		8			
AI 26	6E-05	CC 2			Tm 171		8			
CI 36	2E-04	CC 2			Lu 174		8			
Ar 39		8			Lu 176		8			
Ar 42		8			Hf 178n		8			
K 40		8			Hf 182		8			
Ca 41	<3E-05	C 3			Pt 193		8			
Mn 53		8			TI 204		8			
Mn 54		8			Pb 205		8			
Fe 55	2.19E-06	CC 2			Pb 210		8			
Co 60	<5.59E-03	C 3			Bi 208		8			
Ni 59	1E-02	CC 2			Bi 210m		8			
Ni 63	9.00E-01	CC 2			Po 210		8			
Zn 65		8			Ra 223		8			
Se 79	3.45E-08	CC 2			Ra 225		8			
Kr 81		8			Ra 226		8			
Kr 85		8			Ra 228		8			
Rb 87		8			Ac 227		8			
Sr 90	2.10E-02	CC 2			Th 227		8			
Zr 93	2E-06	CC 2			Th 228		8			
Nb 91		8			Th 229		8			
Nb 92		8			Th 230		8			
Nb 93m	9.62E-07	CC 2			Th 232		8			
Nb 94		8			Th 234	3E-06	CC 2			
Mo 93		8			Pa 231		8			
Tc 97		8			Pa 233	1.10E-07	CC 2			
Tc 99	1E-05	CC 2			U 232		8			
Ru 106		8			U 233		8			
Pd 107		8			U 234	2.01E-06	CC 2			
Ag 108m	3.90E-06	CC 2			U 235	6E-08	CC 2			
Ag 110m		8			U 236	3.00E-07	CC 2			
Cd 109		8			U 238	3E-06	CC 2			
Cd 113m	<9.45E-05	C 3			Np 237	1.10E-07	CC 2			
Sn 119m		8			Pu 236		8			
Sn 121m	<3.28E-04	C 3			Pu 238	3.56E-04	CC 2			
Sn 123		8			Pu 239	9.00E-04	CC 2			
Sn 126	3.04E-07	CC 2			Pu 240	9.99E-04	CC 2			
Sb 125	9.10E-08	CC 2			Pu 241	9.74E-03	CC 2			
Sb 126	4.26E-08	CC 2			Pu 242	6E-07	CC 2			
Te 125m	2.28E-08	CC 2			Am 241	2.30E-03	CC 2			
Te 127m		8			Am 242m	3.71E-06	CC 2			
l 129	2E-08	CC 2			Am 243	9.99E-07	CC 2			
Cs 134	1.30E-08	CC 2			Cm 242	3.06E-06	CC 2			
Cs 135	7E-07	CC 2			Cm 243	4.95E-07	CC 2			
Cs 137	2.83E-02	CC 2			Cm 244	5.63E-06	CC 2			
Ba 133	<3.36E-05	C 3			Cm 245		8			
La 137		8			Cm 246		8			
La 138		8			Cm 248		8			
Ce 144		8			Cf 249		8			
Pm 145		8			Cf 250		8			
Pm 147	1.52E-06	CC 2			Cf 251		8			
Sm 147		8			Cf 252		8			
Sm 151	1.78E-04	CC 2			Other a					
Eu 152	1.39E-06	CC 2			Other b/g					
Eu 154	5.95E-05	CC 2			Total a	4.57E-03	CC 2	0		
Eu 155	7.16E-06	CC 2			Total b/g	9.85E-01	CC 2	0		
	<u>I</u>					<u>i </u>		<u>i </u>		

Bands (Upper and Lower)

A a factor of 1.5 B a factor of 3 C a factor of 10 D a factor of 100

D a factor of 100 E a factor of 1000

Note: Bands quantify uncertainty in mean radioactivity.

Code

- 1 Measured activity
- 2 Derived activity (best estimate)
- 3 Derived activity (upper limit)
- 4 Not present
- 5 Present but not significant
- 6 Likely to be present but not assessed
- 7 Present in significant quantities but not determined
- 8 Not expected to be present in significant quantity