SITE Berkelev

SITE OWNER **Nuclear Decommissioning Authority**

No

WASTE CUSTODIAN Magnox Limited

ILW WASTE TYPE

Is the waste subject to

WASTE VOLUMES

Scottish Policy:

Reported

Stocks: At 1.4.2022..... 27.5 m³

Total future arisings: $0 \, \text{m}^3$

Total waste volume: 27.5 m³

Stock (upper):

Comment on volumes: The last arising of this waste was in 1982. There will be no further arisings of this waste

x 1.1

x 0.9

stream

Uncertainty factors on

volumes: Stock (lower): Arisings (upper)

Arisings (lower) х

WASTE SOURCE The waste is miscellaneous contaminated items arising at Berkeley Nuclear Laboratories

(now Berkeley Technology Centre), principally from the examination of irradiated fuel, steel

and graphite.

PHYSICAL CHARACTERISTICS

General description: The waste consists principally of non-fuel element debris from the post irradiation

> examination work carried out on fuel, steel and graphite in the BC caves and cells. This waste will also comprise combustible and non-combustible wastes, metals, glass and cave refurbishment wastes. There will be a small amount of fire suppressant powders and vermiculite. The waste is contained within mild steel black can liners and paint tin liners. Current plans are that the mild steel waste containers described in former stream 9A926 will not be separated from the waste in this stream. Originally, this waste stream included part of the 3m³ of dried ion exchange material that was used as an infill for approximately 175 of the MCI black cans between 1975 and 1987. This mobile waste has now got a separate identifier (9A82) and therefore, the volumes for 9A69 and 9A70 have been reduced by 1.5m³ each. As the waste is containerised, it is unlikely that there will be any

large items that will require special handling.

The waste will include steel items such as tools, plastic items, tissues and swabs used for Physical components (%vol):

cleaning, fire suppressant powders and the absorber vermiculite. There may be small quantities of ion exchange material, but this is described in stream 9A82 and will be treated with stream 9A25. % Breakdown, ~25% ferrous metals, ~25% glass, ~20% plastics, ~15% paper, ~ 5% each of organic and inorganic Ion Exchange resins and ~5% of ternary

eutectic chloride.

Sealed sources: The waste does not contain sealed sources.

Bulk density (t/m3): 0.5

Comment on density: Densities calculated using weights and external volumes of containers.

CHEMICAL COMPOSITION

General description and components (%wt):

The waste will include steel, plastics, paper and graphite. There may be small amounts of ternary eutectic chloride (sodium chloride, barium chloride and potassium chloride) which was used as fire extinguishant, and incinerator ash. Fission products, actinides and other

activation products will be present as contaminants.

Chemical state: Alkali

Chemical form of radionuclides:

H-3: Most tritium is expected to be present as water but some may be in the form of other

inorganic compounds or as organic compounds.

C-14: Chemical form of carbon 14 has not been determined but may be graphite.

CI-36: The chemical form of chlorine 36 in these wastes is not known.

U: Chemical form of uranium isotopes has not been determined but may be uranium

oxides.

Pu: Chemical form of plutonium isotopes has not been determined but may be plutonium

Metals and alloys (%wt): Much of the metal will be of only 1-2 mm thickness but there will be items of greater

thickness.

| | (%wt) | Type(s) / Grade(s) with proportions | % of total C14 activity |
|----------------------|-------|--|-------------------------|
| Stainless steel | NE | | - |
| Other ferrous metals | ~25.0 | | |
| Iron | | | |
| Aluminium | NE | | |
| Beryllium | NE | | |
| Cobalt | | | |
| Copper | NE | | |
| Lead | NE | | |
| Magnox/Magnesium | NE | | |
| Nickel | | | |
| Titanium | | | |
| Uranium | | | |
| Zinc | NE | | |
| Zircaloy/Zirconium | NE | | |
| Other metals | NE | "Other" metals have not been identified. | |

Organics (%wt):

There will be a number of organic materials present in the waste, including plastics, Perspex, polythene, PVC and paper. Specific organic chemical compounds identified as possibly present are: hexamine, methanol, Perklone D, acetone and epoxy resin. These compounds may adversely affect radionuclide mobility. No quantitative estimates of the various types or quantities of organics have been assessed. There may also be small quantities of liquid organic oils, e.g. oil on swabs and in hydraulic equipment.

| | (%wt) | Type(s) and comment | % of total C14 activity |
|--------------------------------|-------|---------------------|-------------------------|
| Total cellulosics | ~15.0 | | activity |
| Paper, cotton | ~15.0 | | |
| Wood | NE | | |
| Halogenated plastics | NE | | |
| Total non-halogenated plastics | NE | | |
| Condensation polymers | NE | | |
| Others | NE | | |
| Organic ion exchange materials | ~5.0 | | |
| Total rubber | NE | | |
| Halogenated rubber | NE | | |
| Non-halogenated rubber | NE | | |
| Hydrocarbons | | | |
| Oil or grease | | | |
| Fuel | | | |
| Asphalt/Tarmac (cont.coal tar) | | | |
| Asphalt/Tarmac (no coal tar) | | | |
| Bitumen | | | |
| Others | | | |
| Other organics | NE | | |

Other materials (%wt):

| Other materials (%wt): - | | |
|--|------------|---|
| | (%wt) | Type(s) and comment % of total C14 activity |
| Inorganic ion exchange materials | ~5.0 | |
| Inorganic sludges and flocs | 0 | |
| Soil | 0 | |
| Brick/Stone/Rubble | 0 | |
| Cementitious material | 0 | |
| Sand | | |
| Glass/Ceramics | ~25.0 | |
| Graphite | NE | |
| Desiccants/Catalysts | | |
| Asbestos | 0 | |
| Non/low friable | | |
| Moderately friable | | |
| Highly friable | | |
| Free aqueous liquids | 0 | |
| Free non-aqueous liquids | NE | |
| Powder/Ash | NE | |
| Inorganic anions (%wt): The inorganic anion | content of | f the waste has not been quantified. |
| | (%wt) | Type(s) and comment |
| Fluoride | 0 | |
| Chloride | ~5.0 | |
| lodide | 0 | |
| Cyanide | 0 | |
| Carbonate | NE | |
| Nitrate | NE | |
| Nitrite | NE | |
| Phosphate | NE | |
| Sulphate | NE | |
| Sulphide | 0 | |
| waste acceptance criteria: corrosive effect of s | odium chlo | c) fire extinguishant is of potential concern owing to the oride (one third of total contents of TEC) on the waste chromium trioxide has been identified as possibly present. |
| | (%wt) | Type(s) and comment |
| Combustible metals | TR | |
| Low flash point liquids | 0 | |
| | 0 | |

| | (%wt) | Type(s) and comme |
|---------------------------|-------|-------------------|
| Combustible metals | TR | |
| Low flash point liquids | 0 | |
| Explosive materials | 0 | |
| Phosphorus | 0 | |
| Hydrides | TR | |
| Biological etc. materials | 0 | |
| Biodegradable materials | 0 | |
| Putrescible wastes | 0 | |
| | | |

2022 Inventory

| Non-putrescible wastes | | |
|--|---------------------------|--|
| Corrosive materials | 0 | |
| Pyrophoric materials | 0 | |
| Generating toxic gases | 0 | |
| Reacting with water | TR | |
| Higher activity particles | | |
| Soluble solids as bulk chemical compounds | | |
| Hazardous substances / None expected non hazardous pollutants: | d, | |
| | (%wt) Type(s) and comment | |
| Acrylamide | | |
| Benzene | | |
| Chlorinated solvents | | |
| Formaldehyde | | |
| Organometallics | | |
| Phenol | | |
| Styrene | | |
| Tri-butyl phosphate | | |
| Other organophosphates | | |
| Vinyl chloride | | |
| Arsenic | | |
| Barium | | |
| Boron | 0 | |
| Boron (in Boral) | | |
| Boron (non-Boral) | | |
| Cadmium | | |
| Caesium | | |
| Selenium | | |
| Chromium | | |
| Molybdenum | | |
| Thallium | | |
| Tin | | |
| Vanadium | | |
| Mercury compounds | | |
| Others | | |
| Electronic Electrical Equipment | (EEE) | |
| EEE Type 1 | | |
| EEE Type 2 | | |
| EEE Type 3 | | |
| EEE Type 4 | | |
| EEE Type 5 | | |

| Complexing a | gents (%wt): | Yes | | | | | |
|----------------------------------|------------------------------------|--|-------------|------------------------|-------------------------|-----------------|--------------------|
| | | | (%wt) | Type(s) | and comment | | |
| I | EDTA | | | | | | |
| i | DPTA | | | | | | |
| 1 | NTA | | | | | | |
| I | Polycarboxylic ad | oids | | | | | |
| (| Other organic co | mplexants | TR | Organic small qu | complexing ag antities. | ents may be p | present in |
| - | Total complexing | agents | ~1.0 | | | | |
| Potential for th contain discret | | Yes. In & of itself no steel components) | ot a DI; wa | iste streai | m may include | DIs (notably a | any stainless |
| PACKAGING | AND CONDIT | TONING | | | | | |
| Conditioning m | nethod: | This stream is to be c 9A68, 9A70, 9A71, 9A 9A68, 9A71 & 9A75. | | | | | |
| Plant Name: | | - | | | | | |
| Location: | | Berkeley Site | | | | | |
| Plant startup d | ate: | - | | | | | |
| Total capacity (m³/y incoming | waste): | - | | | | | |
| Target start da packaging this | | - | | | | | |
| Throughput for (m³/y incoming | | - | | | | | |
| Other informat | ion: | - | | | | | |
| Likely containe type: | Container | | pacl | aste kaged %vol) | Waste loading (m³) | Payload (m³) | Number of packages |
| | | | | | | | |
| | | | . | | | | |
| Likely containe comment: | er type | - | | | | | |
| Range in contavolume: | Range in container waste . volume: | | | | | | |
| Other informat containers: | ion on | - | | | | | |
| Likely condition | ning matrix: | | | | | | |

Other information:

Conditioned density (t/m³): Conditioned density comment:

Other informs

Other information on conditioning:

Opportunities for alternative disposal routing:

2022 Inventory

Estimated Date that Opportunity Comment Opportunity Confidence volume (%)

will be realised

Stream

RADIOACTIVITY

Baseline

Source: The waste has become contaminated from the processes concerned with the examination

of irradiated fuel, steel and graphite at Berkeley Nuclear Laboratories (now Berkeley

Technology Centre).

Opportunity

Management Route Management Route

Uncertainty: The values quoted are indicative of the activities that might be expected.

Definition of total alpha and total beta/gamma:

Where totals are shown on the table of radionuclide activities they are the sums of the listed alpha or beta/gamma emitting radionuclides plus 'other alpha' or 'other beta/gamma'.

Measurement of radioactivities:

The specific activities were derived by estimation based upon available information.

Other information: Specific activity is a function of operating history.

| | Mean radioactivity, TBq/m³ | | | Mean radioactivity, TBq/m³ | | | | | |
|-------------------|----------------------------|-----------|----------|----------------------------|-------------------|-------------------|--------------|----------|-----------|
| Nuclide | Waste at | Bands and | Future | Bands and | Nuclide | Waste at | Bands and | Future | Bands and |
| | 1.4.2022 | Code | arisings | Code | | 1.4.2022 | Code | arisings | Code |
| H 3 Be 10 | 8.63E-04 | CC 2 8 | | | Gd 153 Ho 163 | | 8 8 | | |
| C 14 | 9.99E-06 | CC 2 | | | Ho 166m | | 8 | | |
| Na 22 | 9.99E-06 | 8 | | | Tm 170 | | 8 | | |
| Al 26 | | 8 | | | Tm 170 | | 8 | | |
| CI 36 | 7E-07 | CC 2 | | | Lu 174 | | 8 | | |
| Ar 39 | 72-07 | 8 | | | Lu 174 | | 8 | | |
| Ar 42 | | 8 | | | Hf 178n | | 8 | | |
| K 40 | | 8 | | | Hf 182 | | 8 | | |
| Ca 41 | | 8 | | | Pt 193 | | 8 | | |
| Mn 53 | | 8 | | | TI 204 | | 8 | | |
| Mn 54 | | 8 | | | Pb 205 | | 8 | | |
| Fe 55 | 4.39E-06 | CC 2 | | | Pb 210 | | 8 | | |
| Co 60 | 4.17E-05 | CC 2 | | | Bi 208 | | 8 | | |
| Ni 59 | 1E-06 | CC 2 | | | Bi 210m | | 8 | | |
| Ni 63 | 7.21E-05 | CC 2 | | | Po 210 | | 8 | | |
| Zn 65 | | 8 | | | Ra 223 | | 8 | | |
| Se 79 | 1.21E-08 | CC 2 | | | Ra 225 | | 8 | | |
| Kr 81 | | 8 | | | Ra 226 | | 8 | | |
| Kr 85 | | 8 | | | Ra 228 | | 8 | | |
| Rb 87 | | 8 | | | Ac 227 | | 8 | | |
| Sr 90 | 4.89E-03 | CC 2 | | | Th 227 | | 8 | | |
| Zr 93 | 6E-07 | CC 2 | | | Th 228 | | 8 | | |
| Nb 91 Nb 92 | | 8 | | | Th 229 Th 230 | | 8 8 | | |
| Nb 93m | 3.85E-07 | 8 CC 2 | | | Th 232 | | 8 | | |
| Nb 94 | 3.83L-07 | 8 | | | Th 234 | 3E-07 | CC 2 | | |
| Mo 93 | | 8 | | | Pa 231 | 3L-01 | 8 | | |
| Tc 97 | | 8 | | | Pa 233 | 4.16E-08 | CC 2 | | |
| Tc 99 | 3E-06 | CC 2 | | | U 232 | | 8 | | |
| Ru 106 | | 8 | | | U 233 | | 8 | | |
| Pd 107 | | 8 | | | U 234 | 3.09E-07 | CC 2 | | |
| Ag 108m | <2.94E-06 | C 3 | | | U 235 | 7E-09 | CC 2 | | |
| Ag 110m | | 8 | | | U 236 | 4.00E-08 | CC 2 | | |
| Cd 109 | | 8 | | | U 238 | 3E-07 | CC 2 | | |
| Cd 113m | | 8 | | | Np 237 | 4.16E-08 | CC 2 | | |
| Sn 119m | | 8 | | | Pu 236 | | 8 | | |
| Sn 121m | | 8 | | | Pu 238 | 1.78E-04 | CC 2 | | |
| Sn 123 | 4.055.00 | 8 | | | Pu 239 | 1.00E-04 | CC 2 | | |
| Sn 126 | 4.35E-08 | CC 2 | | | Pu 240 | 2.00E-04 | CC 2 | | |
| Sb 125 | 6.09E-09 | 8 CC 2 | | | Pu 241 | 1.46E-03 1E-07 | CC 2 CC 2 | | |
| Sb 126 Te 125m | 0.09E-09 | 8 | | | Pu 242 Am 241 | 1E-07 3.44E-04 | CC 2 | | |
| Te 127m | | 8 | | | Am 242m | 8.36E-07 | CC 2 | | |
| I 129 | 6E-09 | CC 2 | | | Am 243 | 3.00E-07 | CC 2 | | |
| Cs 134 | | 8 | | | Cm 242 | 6.90E-07 | CC 2 | | |
| Cs 135 | 1E-07 | CC 2 | | | Cm 243 | 1.41E-07 | CC 2 | | |
| Cs 137 | 4.95E-03 | CC 2 | | | Cm 244 | 1.69E-06 | CC 2 | | |
| Ba 133 | | 8 | | | Cm 245 | | 8 | | |
| La 137 | | 8 | | | Cm 246 | | 8 | | |
| La 138 | | 8 | | | Cm 248 | | 8 | | |
| Ce 144 | | 8 | | | Cf 249 | | 8 | | |
| Pm 145 | 0.405.00 | 8 | | | Cf 250 | | 8 | | |
| Pm 147 | 9.46E-08 | CC 2 | | | Cf 251 | | 8 | | |
| Sm 147 | 1 795 05 | 8 CC 2 | | | Cf 252 Other a | | 8 | | |
| Sm 151 Eu 152 | 1.78E-05 1.39E-07 | CC 2 | | | Other b/g | | | | |
| Eu 154 | 8.93E-06 | CC 2 | | | Total a | 8.25E-04 | CC 2 | 0 | |
| Eu 155 | 3.59E-07 | CC 2 | | | Total b/g | 1.23E-02 | CC 2 | 0 | |
| | I -::-52 5, | | | | | 1 | | <u> </u> | |

Bands (Upper and Lower)

A a factor of 1.5 B a factor of 3 C a factor of 10 D a factor of 100

E a factor of 1000

Note: Bands quantify uncertainty in mean radioactivity.

Code

- 1 Measured activity
- 2 Derived activity (best estimate)
- 3 Derived activity (upper limit)
- 4 Not present
- 5 Present but not significant
- 6 Likely to be present but not assessed
- 7 Present in significant quantities but not determined 8 Not expected to be present in significant quantity